



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Protocol for a Randomized Sham-Controlled Double-blind Multicenter Efficacy Study of the Gelstix™ Nucleus Augmentation Device to treat Chronic Discogenic Low Back Pain

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-053772
Article Type:	Protocol
Date Submitted by the Author:	28-May-2021
Complete List of Authors:	<p>Maino, P.; Neurocenter of Southern Switzerland, EOC, Pain Management Center; Università della Svizzera Italiana, Faculty of Biomedical Sciences</p> <p>Kallewaard, Jan-Willem; Rijnstate, Anesthesiology</p> <p>van Kuijk, Sander; Maastricht Universitair Medisch Centrum+, Clinical Epidemiology and Medical Technology Assessment</p> <p>Dukanac, J.; Neurocenter of Southern Switzerland, EOC, Pain Management Center</p> <p>Scascighini, L.; University of Applied Sciences and Arts of Southern Switzerland, Department of Health Sciences</p> <p>Cianfoni, A.; Neurocenter of Southern Switzerland, EOC, Service of Diagnostic and Interventional Neuroradiology; Inselspital University Hospital Bern, Dept. of Neuroradiology</p> <p>Scarone, P.; Neurocenter of Southern Switzerland, EOC, Clinic of Neurosurgery</p> <p>Kuhlen, D.E.; Neurocenter of Southern Switzerland, EOC, Clinic of Neurosurgery</p> <p>Hollman, M; Amsterdam UMC Locatie AMC, Department of Anesthesiology</p> <p>Koetsier, Eva; Neurocenter of Southern Switzerland, EOC, Pain Management Center; Università della Svizzera Italiana, Faculty of Biomedical Sciences</p>
Keywords:	Back pain < ORTHOPAEDIC & TRAUMA SURGERY, PAIN MANAGEMENT, Musculoskeletal disorders < ORTHOPAEDIC & TRAUMA SURGERY

SCHOLARONE™  
Manuscripts

1

2

31 Randomized Sham-Controlled Double-blind Multicenter Efficacy Study of the Gelstix™ Nucleus

4

52 Augmentation Device to treat Chronic Discogenic Low Back Pain

6

73

8

9

104 P. Maino<sup>1,2\*</sup>, J.W. Kallewaard<sup>3,4\*</sup>, S.M.J. van Kuijk<sup>5</sup>, J. Dukanac<sup>1</sup>, L. Scascighini<sup>6</sup>, A. Cianfoni<sup>7,8</sup>, P.

11

125 Scarone<sup>9</sup>, D.E. Kuhlen<sup>9</sup>, M.W. Hollmann<sup>4</sup>, E. Koetsier<sup>1,2</sup>

13

146

15

167 \* Both authors contributed equally and therefore share first authorship

17

188

19

209 1. Pain Management Center, Neurocenter of Southern Switzerland, EOC, Lugano, Switzerland

21

2210 2. Faculty of Biomedical Sciences, Università della Svizzera Italiana, Lugano, Switzerland

23

2411 3. Department of Anesthesiology and Pain Management Arnhem, Rijnstate Hospital, Velp, the

25

2612 Netherlands

27

2813 4. Department of Anesthesiology, Amsterdam UMC, location Academic Medical Center, Amsterdam,

29

3014 Netherlands

31

3215 5. Department of Clinical Epidemiology and Medical Technology Assessment, Maastricht University

33

3416 Medical Centre+, Maastricht, the Netherlands

35

3617 6. Department of Health Sciences, University of Applied Sciences and Arts of Southern Switzerland,

37

3818 Manno, Switzerland

39

4019 7. Service of Diagnostic and Interventional Neuroradiology, Neurocenter of Southern Switzerland,

41

4220 EOC, Lugano, Switzerland

43

4421 8. Dept. of Neuroradiology, Inselspital University Hospital Bern, Switzerland

45

4622 9. Clinic of Neurosurgery, Neurocenter of Southern Switzerland, EOC, Lugano, Switzerland

47

4823

49

5024 Word count: 3840

51

5225 Number of tables: 0

53

5426 Number of figures: 3

55

56

57

58

59

60

1  
2  
3 1  
4  
5 2 Address correspondence and reprint requests to:  
6  
7 3 Dr. Eva Koetsier, Pain Management Center, Neurocenter of Southern Switzerland, EOC, Lugano,  
8  
9 4 Switzerland.  
10  
11 5 Ospedale Regionale di Lugano, Sede Ospedale Italiano, via Pietro Capelli 1, 6962, business telephone  
12  
13 6 number: +41 (0)91 8117590, fax number: +41 (0)91 8117591, E-mail: [evakoetsier6@gmail.com](mailto:evakoetsier6@gmail.com),  
14  
15 7 [eva.koetsier@eoc.ch](mailto:eva.koetsier@eoc.ch).  
16  
17 8  
18  
19 9 Sponsor-Investigator contact information:  
20  
21 10 Dr. Paolo Maino, Pain Management Center, Neurocenter of Southern Switzerland, EOC, Lugano,  
22  
23 11 Switzerland.  
24  
25 12 Ospedale Regionale di Lugano, Sede Ospedale Italiano, via Pietro Capelli 1, 6962, Lugano,  
26  
27 13 Switzerland, business telephone number: +41 (0)91 8117590, fax number: +41 (0)91 8117591, E-mail:  
28  
29 14 [paolo.maino@eoc.ch](mailto:paolo.maino@eoc.ch).  
30  
31  
32  
33  
34 15  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1

2

31 ABSTRACT

4

52 Introduction

6

73 Discogenic pain is the cause of pain in up to 40% of patients consulting a physician for low back pain.

8

94 Consensus about treatment of chronic discogenic low back pain is lacking and the majority of

10

115 treatment alternatives is supported by limited evidence, and fusion surgery is not proven to be

12

136 superior to conservative treatment. We hypothesize that treatment with GelStix™ will lead to greater

14

157 reduction in pain intensity at six months post-treatment compared to patients receiving sham

16

178 treatment.

18

199 Methods and analysis

20

2110 This is a parallel group, randomized sham-controlled double-blind, multicentre trial to assess whether

22

2311 the GelStix™ device is superior to sham in reducing pain intensity in patients with chronic discogenic

24

2512 low back pain. The primary outcome will be the change in pain intensity between preoperative baseline

26

2713 and at six months post-intervention. Secondary outcomes include disability, quality of life, the patient's

28

2914 global impression of change scale, the use of pain medication, and the disc degeneration process

30

3115 assessed by means of MRI. For change in pain intensity, disability, health related quality of life, and

32

3316 disc height, mean values will be compared between groups using linear regression analysis, adjusted

34

3517 for treatment centre.

36

3718 Ethics and dissemination

38

3919 Ethics approval was obtained from the Ethics Committee of the Canton Ticino, Switzerland (CE2982)

40

4120 and by the Medical Ethical Committee Arnhem-Nijmegen, the Netherlands (2016-2944). Results will

42

4321 be disseminated through international publications in peer reviewed journals, in addition to

44

4522 international conference presentations.

46

4723

48

4924 Trial registration number NCT02763956

50

51

5225 Protocol version 7.1, 18/11/2020

53

54

5526 Keywords Back pain, pain management, musculoskeletal disorders

56

57

58

59

60

### ARTICLE SUMMARY

#### Strengths and limitations of this study

► This will be the first prospective, randomized, controlled, multicentre trial assessing effectivity and safety of the GelStix™ Nucleus Augmentation Device compared to a sham control in patients with lumbar discogenic pain that had no benefit from conservative care.

► Means to reduce risk of bias are implemented, which includes an a-priori sample size calculation, an explicitly stated primary hypothesis to be tested, methodological rigor, double-blinding, randomization, adequate concealment of group allocation and the assessment of the success of blinding in participants and observers.

► This is also the first study that assesses the disc degeneration process and disc height by means of Magnetic Resonance Imaging (MRI) one year after GelStix™ implantation versus sham.

► All participants will also be treated according to a protocolized physiotherapy.

► The limitations are those inherent to a prospective, randomized sham-controlled double-blind study, including strict exclusion criteria and thus limited generalizability (e.g., protrusions in contact with any nerve root at the symptomatic level or >5mm, an insufficient number of patients, and adherence to a strict protocol that does not necessarily reflect real word daily practice.

1

2

31 INTRODUCTION

4

52 Background and rationale

6

7

83 Discogenic low back pain is characterized by persistent, predominantly centralized axial low back

9

104 pain that worsens with axial loading. It is associated with intervertebral disc degeneration without

11

125 herniation,<sup>1–4</sup> and is thought to be the cause of pain in up to 40% of patients consulting a physician

13

146 for low back pain.<sup>5–8</sup> The water-binding capabilities of the intervertebral disc diminish with aging<sup>9</sup>

15

167 leading to progressive shrinking of the nucleus pulposus and loss of elasticity.<sup>9–12</sup> The cartilaginous

17

188 endplate vascular flow decreases due to a progressive loss in vascularization leading to accumulation

19

209 of cellular waste products, and an increasingly acidic environment.<sup>9,13</sup> A low pH around the discus is

21

2210 associated with discogenic pain.<sup>14,15</sup>

23

2411 Medical history, physical examination, and imaging (e.g. magnetic resonance imaging (MRI)) provide

25

2612 inadequate sensitivity and specificity to accurately diagnose discogenic pain.<sup>16,17</sup> Despite an ongoing

27

2813 debate, moderate evidence supports diagnostic accuracy of provocative discography.<sup>18–20</sup> While

29

3014 previous studies suggest that high-pressure provocative discography may accelerate disc

31

3215 degeneration,<sup>21–23</sup> a recently published study suggests that low-pressure provocative discography,

33

3416 performed according to International Association for the Study of Pain (IASP) criteria, does not

35

3617 accelerate disc degeneration.<sup>24</sup>

37

3818 Consensus about treatment of chronic discogenic low back pain is lacking and the majority of

39

4019 treatment alternatives is supported by limited evidence.<sup>1,4</sup> Conservative management includes anti-

41

4220 inflammatory drugs, physiotherapy, and multidisciplinary biopsychosocial rehabilitation.<sup>25</sup> If

43

4421 conservative treatment fails, (minimally) invasive treatments are considered. Most minimally

45

4622 invasive treatments, such as intradiscal injections (e.g. with methylene blue) and thermal

47

4823 intradiscal/annular techniques (intradiscal electrothermal therapy (IDET), have been abandoned

49

5024 because of poor evidence.<sup>26–28</sup> A recent systematic review concluded that most minimal invasive

51

5225 treatments for discogenic low back have very low evidence; only biacuplasty has moderate evidence

53

5426 for a subgroup of patients with discogenic low back pain.<sup>29</sup>

55

56

57

58

59

60

1 Fusion surgery and total disc replacement, although contemplated as possible therapies in some  
2 cases, are invasive interventions associated with risk of adjacent segment disorder and morbidity.<sup>4,30</sup>  
3 In addition, fusion surgery is not superior to conservative treatment with multidisciplinary  
4 biopsychosocial rehabilitation and physiotherapy.<sup>31,32</sup> Recently, with the emergence of new  
5 frequencies (burst, dorsal root ganglion stimulation, high frequency-10Hz), low back pain has  
6 become a good treatment option for neuromodulation. Considering the fact that neuromodulation is  
7 a more invasive treatment the need is great to find evidence for minimal invasive treatment for  
8 chronic discogenic low back pain.<sup>33,34</sup>  
9 Therefore, treatment options filling the gap between conservative care and invasive surgical  
10 intervention are urgently needed. Currently the first studies are published showing effect of the use  
11 of platelet-rich plasma (PRP) and mesenchymal signaling cells (MSCs) for discogenic pain. Notably, no  
12 intervention has multiple RCT's published yet.<sup>35</sup> The implantation of hydrogels into the nucleus  
13 pulposus represents a promising regenerative intradiscal therapy, in particular in patients with early  
14 or moderate disc degeneration not responding to conservative care.<sup>36,37</sup> The hydrogel containing  
15 'GelStix™ Nucleus Augmentation Device' (hereafter called GelStix™) is composed primarily of  
16 hydrolyzed polyacrylonitrile (HPAN). The GelStix™ is shaped in the form of an elongated matchstick  
17 and can be inserted percutaneously into the nucleus through a needle. Once implanted, the GelStix™  
18 absorbs the body's own fluids and expands around tenfold in volume (see Fig. 1).

19  
20 **Insert here Figure 1**

21  
22 The GelStix™ material acts as a reservoir of permanent hydration of the intervertebral disc,  
23 producing increased pressure, and improved fluid exchange and pH balance, leading to disc  
24 preservation.<sup>38</sup> Results of previous non-controlled studies suggest that GelStix™ implantation leads to  
25 a significant pain and disability relief four weeks after implantation in patients with discogenic  
26 pain.<sup>39,40</sup>





## METHODS AND ANALYSIS

This protocol has been written in accordance with the Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT) checklist. The study will be conducted in two regional hospitals: the Pain Management Center, Neurocenter of Southern Switzerland, Lugano, Switzerland, and the Department of Anaesthesiology and Pain Management Arnhem, Rijnstate Hospital, Arnhem, the Netherlands.

### Participants

The target population is represented by patients suffering from discogenic low back pain with a baseline numeric rating scale (NRS) pain score  $\geq 5/10$  following at least twelve weeks conservative care.

#### Inclusion criteria:

- 18-66 years of age
- Lumbar DDD on MRI scan with Pfirrmann grade<sup>41</sup> 2, 3 or 4
- Discogenic pain confirmed by positive discography\* of one or maximum two lumbar disc levels, and one negative control level
- Persistent predominant, nociceptive low back pain with a NRS score of  $\geq 5/10$ , that worsens with axial loading and improves with recumbence of at least 12 weeks duration
- Failure to have symptoms resolved or reduced following at least 12 weeks conservative care (drug therapy and/or physiotherapy)
- Negative medial branches block results
- Legally competent and able to understand the nature, scope and aim of the clinical investigation

#### Exclusion criteria:

- Provocative discography will be performed by an experienced pain physician under strict sterile conditions. Thirty minutes before the intervention, intravenous antibiotics for prophylaxis will be administered. The patient will be positioned in the prone position on an X-ray permeable table. After subcutaneous anaesthetic injection of 2 ml mg of lidocaine 1%, the nucleus will be accessed with the

two-needle technique with a 25-27 Gauge needle through the transforaminal, posterolateral approach, according to the technique described by Kallewaard et al.<sup>3</sup> Fluoroscopy will be used to identify spinal levels, guide the needle, and to confirm final needle position. The following variables will be monitored during the injection of the contrast solution: the opening pressure (the pressure at which contrast is first visible in the disc), the provocation pressure (the pressure greater than the opening pressure at which complaints of pain arise), and the peak pressure or the final pressure at the end of the procedure. Additionally, the total volume of the injected contrast solution, the Adams scale,<sup>42</sup> and the pain score measured by NRS per disc level will be recorded.

The procedure, per level, is continued until:<sup>3</sup>

- Concordant pain is reproduced at a level of  $\geq 7/10$  and/or
- The volume infused reaches 3.0 mL and/or
- The pressure rises to 50 psi above opening pressure

According to the guidelines of the IASP,<sup>43</sup> the symptomatic level and the one adjacent level are examined. A disc is only considered to be positive if concordant pain can be induced at the target level (symptomatic level); with an intensity of this pain of at least NRS 7, reproduced by a pressure of less than 50 psi above opening pressure; and if the control level is negative for provocation of pain. A control disc is considered a critical element for defining a positive discography, as it serves as an internal patient control disc and as a possible indicator of central sensitization.

## Interventions

### The GelStix™ implantation

For each participant, up to two levels will be treated. The CE marked GelStix™ Nucleus Augmentation Device system (STX-1835S, Replication Medical, Inc. – Cranbury, NJ, USA), will be implanted by an experienced pain physician familiar with the transforaminal posterolateral discography approach described above. The GelStix™ insertion will be performed under local anaesthesia with a single

needle technique through the procedure-specific 18 Gauge needle (18GTXX165mm, Replication Medical, Inc. – Cranbury, NJ, USA). Up to three GelStixs will be implanted at each symptomatic disc level. Once the needle tip is located in the centre of the nucleus, the stylet will be removed from the needle. Then, the protective cap is removed from the preloaded GelStix™ holder and the GelStix™ holder is threaded onto the proximal end of the introducer needle. The holder stylet is pushed, driving the GelStix™ completely into the introducer needle. The implant holder will then be removed and the needle stylet ('blunt push rod needle') is driven through the needle and bottomed out to deliver the GelStix™ completely into the nucleus, keeping the needle tip centred in the nucleus (fig. 3-9). The procedure will be repeated to insert additional GelStix™. When resistance rises adding a second or third GelStix™, further insertion is discontinued. At the end of the procedure, the needle will be withdrawn, and a sterile bandage will be applied to the insertion site.

**Insert here Figures 3-9.**

#### The sham intervention

For the sham intervention the symptomatic discs will be injected with 1 ml of saline (NaCl 0.9%). Intradiscal saline injection (1 mL NaCl 0.9% ) is safe<sup>44</sup> and has been used as a control/sham intervention in other randomized controlled <sup>26,45,46</sup>

#### Concomitant treatment

Starting two weeks after the intervention, participants of both study groups will be prescribed physiotherapy according to a study specific protocol. Session frequency will be once a week, for nine weeks. An experienced musculoskeletal physiotherapist will assess the patient before starting the post-intervention protocol, in order to determine the starting level for the exercises. Motor control and stabilization exercises will be instructed to the patients and they will get a leaflet with pictures of the exercises to perform at home/at work. Individual exercises include training of the deep

1 abdominal muscles with the lumbar multifidus and the transversus abdominis. Moreover, to restore  
2 the function of the core muscles, all directions and their muscular chains will be trained. All patients  
3 will be instructed as to how to do exercises at home and will be asked to continue these exercises  
4 three times a week for six months. Continuation or modification of pain medication is permitted  
5 during the study period of twelve months.

## 6 7 **Outcome measures**

8 The primary outcome is the change in pain intensity, assessed by means of a pain diary, between  
9 preoperative baseline and at six months post-intervention in the GelStix<sup>tm</sup>-treated compared to the  
10 sham-treated group. Pain intensity will be assessed employing an 11-point (i.e. 0–10) NRS with 0  
11 meaning 'no pain' and '10' meaning 'worst possible pain'.<sup>47</sup> Three times daily pain scores will be  
12 assessed for five consecutive days around the intended measurement time. The mean NRS scores on  
13 the pain diary will furthermore be measured at one week, and one, three, and twelve months.

14  
15 The secondary outcomes include:

- 16 - Disability, using the Oswestry Disability Index (ODI). The ODI is completed at baseline, and at  
17 three, six and twelve months. The ODI is a self-administered questionnaire, assessing the  
18 patient's level of pain and function during basic activities of daily living such as walking,  
19 personal care, standing, sleeping, etc.<sup>48</sup>
- 20 - Quality of life (QoL), quantified with the European Quality of Life Five Dimension Five Level  
21 Scale (EQ-5D-5L). The EQ-5D-5L will be completed at baseline and at three, six and twelve  
22 months. This questionnaire assesses health related quality of life in terms of five dimensions:  
23 mobility, self-care, usual activities, pain/discomfort, and anxiety/depression.<sup>49</sup> Additionally,  
24 the EuroQol Visual Analogue Scale (EQ VAS) records the respondent's self-rated health on a 20  
25 cm vertical, visual analogue scale with endpoints labelled 'the best health you can imagine'  
26 and 'the worst health you can imagine'.

1  
2  
3 1 - The Patient's Global Impression of Change (PGIC) scale will be measured at three, six and  
4  
5 2 twelve months. This scale assesses the patient's own evaluation of improvement or  
6  
7 3 deterioration over time on a 7-point Likert Scale rated from 'very much improved' to 'very  
8  
9 4 much worse'.  
10  
11  
12 5 - The use of pain medication will be assessed as the intake of analgesics at baseline, at one  
13  
14 6 week, and at one, three, six and twelve months.  
15  
16 7 - The disc degeneration process will be assessed by means of MRI twelve months after  
17  
18 8 treatment compared to baseline. Pfirrmann grade,<sup>41</sup> disc height, and the presence of high  
19  
20 9 intensity zones (HIZ),<sup>50</sup> Modic signs,<sup>51</sup> and Schmorl's nodes<sup>52</sup> will be recorded.  
21  
22  
23 10  
24  
25 11 Additionally, to assess the association between pain catastrophizing, surgical fear, state of  
26  
27 12 depression and long-term outcome the following additional patient-reported outcome measures  
28  
29 13 (PROMs) will be registered at baseline. Pain catastrophizing, defined as an exaggerated negative  
30  
31 14 interpretation of the meaning of pain, will be measured by the Pain Catastrophizing Scale (PCS).  
32  
33 15 Higher pain catastrophizing before intervention are related to lower perceived recovery.<sup>53,54</sup> Surgical  
34  
35 16 fear will be measured by the Surgical Fear Questionnaire (SFQ) as a predictor of physical and  
36  
37 17 emotional recovery.<sup>53</sup> State of depression will be assessed by the Hospital Anxiety and Depression  
38  
39 18 Scale (HADS), a self-administered questionnaire developed to detect states of anxiety and depression  
40  
41 19 in hospital out-patient clinics.<sup>55</sup> Moreover, pain self-efficacy will be assessed employing the Pain Self-  
42  
43 20 Efficacy Questionnaire-I (PSEQ-I). This patient self-reported measurement instrument evaluates pain  
44  
45 21 self-efficacy beliefs,<sup>56</sup> i.e. the degree of confidence a patient has in performing regular daily activities  
46  
47 22 despite of pain. The presence of low levels of pain self-efficacy has been shown to be associated with  
48  
49 23 high levels of disability in patients experiencing pain.<sup>57,58</sup>  
50  
51  
52 24  
53  
54  
55 25 The following additional data will be collected at baseline: sex, age, weight, height, smoking habits,  
56  
57 26 previous treatment of discogenic pain, and neurological examination. Employment status baseline and  
58  
59  
60

1 at six and twelve months will be recorded. The proportion of patients unable to return to work will be  
2 an additional measure of efficacy of the treatment.

3 The success of blinding will be assessed at the end of the trial. Before unblinding, the patients and  
4 the blind observers will be asked to guess the patients' treatment and the answers will be compared  
5 with the actual treatments administered. Successful blinding procedures can reduce bias in clinical  
6 trials.<sup>59,60</sup>

7 The safety outcome of this study is the incidence and severity of complications and adverse events  
8 (AE's) including procedure-related complications at any time point in the study. The main expected  
9 adverse device effects are infection (local or discitis), bleeding, nerve damage and/or limited motion  
10 as a result of the procedure.

### 13 **Sample size**

14 Thirty patients per group will be required to have 80% power to detect a minimally clinically relevant  
15 difference of 1.5 points on the NRS between groups, with an estimated standard deviation (SD) of 2,  
16 and testing with an alpha of 5% (two-tailed). With an expected drop-out rate of 20%, 75 patients will  
17 be randomized.

### 20 **Randomization**

21 The Project Manager of the Clinical Trial Unit of the Ente Ospedaliero Cantonale (CTU-EOC),  
22 Bellinzona, Switzerland, will be in charge for computer generated block randomization lists stratified  
23 by centre (blocks of 4). The Project Manager will act as an independent person, not involved in any  
24 other aspect of the trial except administrative/financial issues. The study is patient- and observer-  
25 blinded, while the physician performing the study intervention will necessarily be aware of the



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 treatment allocation. A web-based access to patient allocation codes will be provided to the  
2 physician in charge for GelStix<sup>tm</sup>/placebo injection. The treating team will be instructed not to  
3 communicate allocation to GelStix<sup>tm</sup> or placebo in any way, both to the patient and to other trial  
4 personnel. The “assessors”, i.e., the investigators in charge for efficacy and safety assessments and  
5 the research nurses that may be in charge for questionnaires collection, and the personnel in charge  
6 of monitoring/data review and analysis will have no access to the randomization lists and will receive  
7 no information about patient treatment for the entire duration of the study. For patients still  
8 experiencing substantial discogenic pain at six months, the code can be broken at their request (after  
9 the assessment of the success of blinding). The patients initially allocated to the control group are  
10 then given the opportunity to cross-over to the GelStix<sup>tm</sup> treatment. Any other code breaks should  
11 occur only in circumstances when knowledge of the actual treatment is absolutely essential for  
12 further management of the patient e.g., in case of important AE’s to ensure the most appropriate  
13 patient management.

16 **Data collection and management**

17 Study data will be collected on a case report form by the research team and will be entered in a  
18 research electronic data capture (REDCap) database.<sup>61</sup> The data will be associated to an unique trial  
19 identification number per patient. The database will be double-checked for missing data and data entry  
20 errors. The data from the REDCap database will be imported automatically in the latest version of R, a  
21 language for statistical computing. All study data will be archived for at least of 15 years after study  
22 termination.

24 **Patient involvement**

25 Patient with discogenic pain were involved at several stages of the trial, including the design and  
26 conduct of the trial. We carefully assessed the burden of the trial interventions on these patients. We

1 will disseminate the main results to trial participants and will seek patient and public involvement in  
2 the development of an appropriate method of dissemination.

3

#### 4 **Statistical methods**

5 Baseline characteristics will be described stratified by treatment allocation as mean and standard  
6 deviation or median and first and third quartile, and as count and percentage, as appropriate. In case  
7 of over 5% of missing data, we will use multiple imputation with fully conditional specification to  
8 impute the dataset. The number of imputations will be set to the percentage of incomplete patients.  
9 All subsequent analyses will be performed according to the intention to treat principle. A “per  
10 protocol” analysis will also be performed, excluding patients who are not evaluable for the primary  
11 endpoint because of dropout (e.g., consent withdrawal before completion of the six months  
12 observation period). Frequency and type of AE’s and complications during the study will be described  
13 in the final report. Dropouts will be replaced up to the number of evaluable patients defined in the  
14 sample size calculation.

15 The primary outcome is change in pain (NRS) at six months compared to baseline. Mean values will  
16 be compared between groups using linear regression analysis, adjusted for treatment centre. In case  
17 of imbalance of baseline characteristics as judged by the trial steering committee, regression  
18 analyses will be further adjusted for potential confounders. Change from baseline in pain at other  
19 follow-up moments and change from baseline in continuous secondary outcome measures (i.e.,  
20 disability (ODI) and health related quality of life [EQ-5D-5L], and disc height) will be analysed in a  
21 similar manner. PGIC scores will be dichotomized by taking “very much improved” and “much  
22 improved” to indicate treatment success. Pfirrmann grade will be dichotomized into grade 1 or 2  
23 versus more than 2. Success rates on the PGIC, dichotomized Pfirrmann grade, and the presence of  
24 HIZ, Modic changes, and Schmorl’s nodes will be compared between groups using logistic regression  
25 analysis adjusted for center.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 Univariable and multivariable logistic regression will be used to quantify crude and adjusted  
2 associations between PCS, SFQ, HADS, and PSEQ-I and treatment success. These analyses will be  
3 considered exploratory. The success of blinding will be assessed using the Sign test, testing whether  
4 the percentage of correct guesses differs from that expected by chance (i.e. 50%).  
5

6 **Monitoring**

7 The research project will be monitored by a certified clinical monitor, which will review the data  
8 quality and will ensure that study activities are carried out in accordance with the protocol, good  
9 clinical practice and applicable regulatory requirements. This being a novel treatment method, a  
10 blinded interim analysis for futility will be planned for the primary outcome measure at T3 months  
11 after 40 patients (i.e. 20 in each arm of the study) have been enrolled. The study will be terminated  
12 in case the experimental arm performs significantly worse (as based on independent samples t-test  
13 or Mann-Whitney-U test) *and* the difference between groups is clinically relevant (i.e. 2 points or  
14 more on the NRS).  
15  
16

17 **Limitations of the study**

18 The limitations are those inherent to a prospective, randomized, sham-controlled study, including  
19 difficulty in recruiting patients due to potential patient refusal and strict exclusion criteria (e.g.,  
20 protrusions in contact with any nerve root on the symptomatic level or >5mm), an insufficient  
21 number of patients, and adherence to a strict protocol that does not necessarily reflect real world  
22 daily practice. Recently performed strategies for achieving adequate participant enrolment to reach  
23 target sample size are the drafting and dispersal of an informative letter to referral colleagues in  
24 Switzerland and in the Netherlands, the introduction of a back pain treatment algorithm in the Pain  
25 Management Center in Lugano, indicating a clear algorithm to follow after negative medial branch  
26 block tests, indicating also the possibility for inclusion in the GelStix™ study.

Another limitation of this trial is the question whether intradiscal saline injection is a true placebo. For example, a recently published systematic review and meta-analysis of Manchikanti et al. showed that epidurally administered saline and saline with steroids may be both effective in managing low back and lower extremity pain.<sup>62</sup> On the other hand, saline has been routinely used as a sham intervention in several other intradiscal treatment studies such as the randomized controlled trial (RCT) of Kallewaard et al.,<sup>26</sup> which compared intradiscal methylene blue plus lidocaine to intradiscal saline plus lidocaine injection, and two the RCT's of Cao et al.<sup>45</sup> and Khot et al.<sup>46</sup> comparing intradiscal corticosteroid to saline injection in the treatment of discogenic low back pain.

1  
2  
3 1 **ETHICS AND DISSEMINATION**  
4

5 2  
6  
7 3 **Research ethics approval and consent to participate**  
8  
9  
10 4 This trial has been approved by the Research Ethics Committee of the Canton Ticino, Switzerland (CE  
11  
12 5 2982) and by the Medical Ethical Committee Arnhem-Nijmegen, the Netherlands (2016-2944). All  
13  
14 6 patients that agree to participate will sign an informed consent form provided by the independent  
15  
16 7 observer. Any amendment to the protocol must as well be approved by this institution.  
17  
18  
19 8  
20  
21 9

22  
23 10 **Confidentiality**  
24

25 11 Individual subject medical information obtained as a result of this study is considered confidential  
26  
27 12 and disclosure to third parties is prohibited. Subject confidentiality will be further ensured by utilizing  
28  
29 13 subject identification code numbers. Direct access to source documents will be permitted for  
30  
31 14 purposes of data review by authorized personnel involved in the trial and inspections. Patients’  
32  
33 15 identity will not be disclosed to the person in charge for the statistical analysis and will not appear in  
34  
35 16 any publication or public presentation of the study results. Results will be disseminated through  
36  
37 17 international publications in peer reviewed journals, in addition to international conference  
38  
39 18 presentations.  
40  
41  
42  
43  
44  
45  
46

47 20 **Funding statement**

48 21 This research received no specific grant from any funding agency in the public, commercial or not-for-  
49  
50 22 profit sectors.  
51

52 23  
53  
54 24 **Declaration of interest**

55  
56 25 The manufacturing Company Replication Medical, Inc. will cover the costs of the GelStix™ material.  
57  
58  
59  
60

1  
2  
3 1 This will be an unrestricted support and the manufacturer will not be involved in study design, data  
4  
5 2 collection, analysis, interpretation and reporting/publication.  
6  
7  
8 3  
9  
10 4  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

References

1. Zhao L, Manchikanti L, Kaye AD, Abd-Elseyed A. Treatment of Discogenic Low Back Pain: Current Treatment Strategies and Future Options—a Literature Review. *Curr Pain Headache Rep*. 2019;23:86.

2. Peng BG. Pathophysiology, diagnosis, and treatment of discogenic low back pain. *World J Orthop*. 2013;4:42-52.

3. Kallewaard JW, Terheggen MAMB, Groen GJ, et al. 15. Discogenic Low Back Pain. *Pain Pract*. 2010;10:560-579.

4. Fujii K, Yamazaki M, Kang JD, et al. Discogenic Back Pain: Literature Review of Definition, Diagnosis, and Treatment. *JBMR Plus*. 2019;3:e10180.

5. Cheung KMC, Karppinen J, Chan D, et al. Prevalence and pattern of lumbar magnetic resonance imaging changes in a population study of one thousand forty-three individuals. *Spine (Phila Pa 1976)*. 2009 Apr 20;34(9):934-940.

6. Schwarzer AC, Aprill CN, Derby R, Fortin J, Kine G. The relative contributions of the disc and zygapophyseal joint in chronic low back pain. *Spine (Phila Pa 1976)*. 1994 Apr 1;19(7):801-806.

7. Verrills P, Nowesenitz G, Barnard A. Prevalence and Characteristics of Discogenic Pain in Tertiary Practice: 223 Consecutive Cases Utilizing Lumbar Discography. *Pain Med (United States)*. 2015;16:1490-1499.

8. DePalma MJ, Ketchum JM, Saullo T. What Is the Source of Chronic Low Back Pain and Does Age Play a Role? *Pain Med*. 2011;12:224-233.

9. Vo NV., Hartman RA, Patil PR, et al. Molecular mechanisms of biological aging in intervertebral discs. *J Orthop Res*. 2016 Aug;34(8):1289-306.

10. Singh K, Masuda K, Thonar EJMA, An HS, Cs-Szabo G. Age-related changes in the extracellular matrix of nucleus pulposus and annulus fibrosus of human intervertebral disc. *Spine (Phila Pa 1976)*. 2009;34:10-16.

11. Patil P, Niedernhofer LJ, Robbins PD, Lee J, Sowa G, Vo N. Cellular Senescence in Intervertebral

- 1 Disc Aging and Degeneration. *Curr Mol Biol Reports*. 2018;4:180-190.
- 2
- 3 12. Kos N, Gradisnik L, Velnar T. A Brief Review of the Degenerative Intervertebral Disc Disease.
- 4
- 5 *Med Arch (Sarajevo, Bosnia Herzegovina)*. 2019;73:421-424.
- 6
- 7 13. Bibby SRS, Jones DA, Ripley RM, Urban JPG. Metabolism of the intervertebral disc: Effects of
- 8
- 9 low levels of oxygen, glucose, and pH on rates of energy metabolism of bovine nucleus
- 10
- 11 pulposus cells. *Spine (Phila Pa 1976)*. 2005;30:487-496.
- 12
- 13 14. Liang CZ, Li H, Tao YQ, et al. The relationship between low pH in intervertebral discs and low
- 14
- 15 back pain: A systematic review. *Arch Med Sci*. 2012;8:952-956.
- 16
- 17 15. Keshari KR, Lotz JC, Link TM, Hu S, Majumdar S, Kurhanewicz J. Lactic acid and proteoglycans as
- 18
- 19 metabolic markers for discogenic back pain. *Spine (Phila Pa 1976)*. 2008;33:312-317.
- 20
- 21 16. Schwarzer AC, Aprill CN, Derby R, Fortin J, Kine G, Bogduk N. The prevalence and clinical
- 22
- 23 features of internal disc disruption in patients with chronic low back pain. *Spine (Phila Pa*
- 24
- 25 *1976)*. 1995;20:1878-1883.
- 26
- 27 17. Brinjikji W, Luetmer PH, Comstock B, et al. Systematic literature review of imaging features of
- 28
- 29 spinal degeneration in asymptomatic populations. *Am J Neuroradiol*. 2015;36:811-816.
- 30
- 31 18. Manchikanti L, Benyamin RM, Singh V, et al. An update of the systematic appraisal of the
- 32
- 33 accuracy and utility of lumbar discography in chronic low back pain. *Pain Physician*. 2013 ;16(2
- 34
- 35 Suppl):SE55-95.
- 36
- 37 19. Manchikanti L, Soin A, Benyamin RM, et al. An update of the systematic appraisal of the
- 38
- 39 accuracy and utility of discography in chronic spinal pain. *Pain Physician*. 2018;21:91-110.
- 40
- 41 20. Wolfer LR, Derby R, Lee JE, Lee SH. Systematic review of lumbar provocation discography in
- 42
- 43 asymptomatic subjects with a meta-analysis of false-positive rates. *Pain Physician*.
- 44
- 45 2008;11:513-538.
- 46
- 47 21. Carragee EJ, Don AS, Hurwitz EL, Cuellar JM, Carrino J, Herzog R. 2009 ISSLS prize winner: Does
- 48
- 49 discography cause accelerated progression of degeneration changes in the lumbar disc: A ten-
- 50
- 51 year matched cohort study. *Spine (Phila Pa 1976)*. 2009;34:2338-2345.
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60



- 1
- 2
- 3 1 22. Cuellar JM, Stauff MP, Herzog RJ, Carrino JA, Baker GA, Carragee EJ. Does provocative
- 4
- 5 2
- 6 discography cause clinically important injury to the lumbar intervertebral disc? A 10-year
- 7
- 8 3
- 9 matched cohort study. *Spine J.* 2016;16:273-280.
- 10 4 23. Koetsier E, Kallewaard JW, Maino P. Commentary on: Does provocative discography cause
- 11
- 12 5
- 13 clinically important injury to the lumbar intervertebral disc? A 10-year matched cohort study,
- 14
- 15 6
- 16 by Cueller et al., *Spine J* 2016. *Spine J.* 2017;17:610-611.
- 17 7 24. McCormick ZL, Lehman VT, Plastaras CT, et al. Low-Pressure Lumbar Provocation Discography
- 18
- 19 8
- 20 According to Spine Intervention Society/International Association for the Study of Pain
- 21
- 22 9
- 23 Standards Does Not Cause Acceleration of Disc Degeneration in Patients with Symptomatic
- 24
- 25 10
- 26 Low Back Pain: A 7-Year Matched Coh. *Spine (Phila Pa 1976).* 2019;44:E1161-E1168.
- 27 11 25. Van Middelkoop M, Rubinstein SM, Kuijpers T, et al. A systematic review on the effectiveness
- 28
- 29 12
- 30 of physical and rehabilitation interventions for chronic non-specific low back pain. *Eur Spine J.*
- 31
- 32 13
- 33 Published online 2011. 2011 Jan;20(1):19-39.
- 34 14 26. Kallewaard JW, Wintraecken VM, Geurts JW, et al. A multicenter randomized controlled trial
- 35
- 36 15
- 37 on the efficacy of intradiscal methylene blue injection for chronic discogenic low back pain:
- 38
- 39 16
- 40 The IMBI study. *Pain.* 2019 Apr;160(4):945-953.
- 41 17 27. Helm S, Hayek SM, Benyamin R, Manchikanti L. Systematic review of the effectiveness of
- 42
- 43 18
- 44 thermal annular procedures in treating discogenic low back pain. *Pain Physician.* 2009;12:207-
- 45
- 46 19
- 47 232.
- 48 20 28. Freeman BJC. IDET: A critical appraisal of the evidence. In: *European Spine Journal.* 2006;15
- 49
- 50 21
- 51 Suppl 3(Suppl 3):S448-457.
- 52 22 29. Huygen F, Kallewaard JW, van Tulder M, et al. "Evidence-Based Interventional Pain Medicine
- 53
- 54 23
- 55 According to Clinical Diagnoses": Update 2018. *Pain Pract.* 2019 Jul;19:664-675.
- 56 24 30. Pan A, Hai Y, Yang J, Zhou L, Chen X, Guo H. Adjacent segment degeneration after lumbar
- 57
- 58 25
- 59 spinal fusion compared with motion-preservation procedures: a meta-analysis. *Eur Spine J.*
- 60
- Published online 2016. 2016;25:1522-1532.

- 1  
2  
3 1 31. Bydon M, De La Garza-Ramos R, Macki M, Baker A, Gokaslan AK, Bydon A. Lumbar fusion  
4  
5 2 versus nonoperative management for treatment of discogenic low back pain: A systematic  
6  
7 3 review and meta-analysis of randomized controlled trials. *J Spinal Disord Tech*. 2014  
8  
9 4 Jul;27(5):297-304.  
10  
11  
12 5 32. Wang X, Wanyan P, Tian JH, Hu L. Meta-analysis of randomized trials comparing fusion  
13  
14 6 surgery to non-surgical treatment for discogenic chronic low back pain. *J Back Musculoskelet*  
15  
16 7 *Rehabil*. 2015;28(4):621-627.  
17  
18 8 33. Vallejo R, Gupta A, Cedeno DL, et al. Clinical Effectiveness and Mechanism of Action of Spinal  
19  
20 9 Cord Stimulation for Treating Chronic Low Back and Lower Extremity Pain: a Systematic  
21  
22 10 Review. *Curr Pain Headache Rep*. 2020;24:70.  
23  
24  
25 11 34. Kallewaard JW, Edelbroek C, Terheggen M, Raza A, Geurts JW. A Prospective Study of Dorsal  
26  
27 12 Root Ganglion Stimulation for Non-Operated Discogenic Low Back Pain. *Neuromodulation*.  
28  
29 13 2020 Feb;23(2):196-202.  
30  
31  
32 14 35. Desai MJ, Mansfield JT, Robinson DM, Miller BC, Borg-Stein J. Regenerative Medicine for Axial  
33  
34 15 and Radicular Spine-Related Pain: A Narrative Review. *Pain Pract*. 2020;20:437-453.  
35  
36 16 36. Gullbrand SE, Schaer TP, Agarwal P, et al. Translation of an injectable triple-interpenetrating-  
37  
38 17 network hydrogel for intervertebral disc regeneration in a goat model. *Acta Biomater*.  
39  
40 18 2017;60:201-209.  
41  
42  
43 19 37. Gan Y, Li P, Wang L, et al. An interpenetrating network-strengthened and toughened hydrogel  
44  
45 20 that supports cell-based nucleus pulposus regeneration. *Biomaterials*. 2017;136:12-28.  
46  
47  
48 21 38. Yue JJ, Morgenstern R, Morgenstern C, Laurysen C. Shape memory hydrogels - A novel  
49  
50 22 material for treating age-related degenerative conditions of the Spine. *Eur Musculoskelet Rev*.  
51  
52 23 2011;6:184-188.  
53  
54 24 39. Sing SH. Lumbar intradiscal treatments: Early evaluation of patient satisfaction in disc repair  
55  
56 25 by nucleus augmentation (GelStix™). *J Tissue Eng Regen Med*. 2012;6 45.  
57  
58  
59 26 40. Ceylan A, Aşık İ, Özgencil GE, Erken B. Clinical results of intradiscal hydrogel administration  
60

- 1
- 2
- 3 1 (GelStix) in lumbar degenerative disc disease. *Turkish J Med Sci.* 2019;49:1634-1639.
- 4
- 5 2 41. Pfirrmann CW, Metzdorf A, Zanetti M, Hodler J, Boos N. Magnetic resonance classification of
- 6
- 7 3 lumbar intervertebral disc degeneration. *Spine (Phila Pa 1976).* 2001;26:1873-1878.
- 8
- 9
- 10 4 42. Adams MA, Dolan P, Hutton WC. The stages of disc degeneration as revealed by discograms. *J*
- 11
- 12 5 *Bone Jt Surg - Ser B.* 1986;68:36-41.
- 13
- 14 6 43. Merskey H, Bogduk N. Classification of Chronic Pain. Descriptions of Chronic Pain Syndromes
- 15
- 16 7 and Definitions of Pain Terms. 2nd ed. Seattle, WA: IASP Press; 1994.
- 17
- 18 8 44. Fukui S, Nitta K, Iwashita N, Tomie H, Nosaka S. Percutaneous intradiscal high-pressure
- 19
- 20 9 injection of saline and lidocaine in patients with lumbar intervertebral disc extrusion. *J Anesth.*
- 21
- 22 10 2012;26:786-789.
- 23
- 24
- 25 11 45. Cao P, Jiang L, Zhuang C, et al. Intradiscal injection therapy for degenerative chronic
- 26
- 27 12 discogenic low back pain with end plate Modic changes. *Spine J.* 2011;11:100-106.
- 28
- 29
- 30 13 46. Khot A, Bowditch M, Powell J, Sharp D. The Use of Intradiscal Steroid Therapy for Lumbar
- 31
- 32 14 Spinal Discogenic Pain: A Randomized Controlled Trial. *Spine (Phila Pa 1976).* 2004;29:833-
- 33
- 34 15 836.
- 35
- 36 16 47. Dworkin RH, Turk DC, Farrar JT, et al. Core outcome measures for chronic pain clinical trials:
- 37
- 38 17 IMMPACT recommendations. *Pain.* 2005;113:9-19.
- 39
- 40
- 41 18 48. Fairbank JCT, Pynsent PB. The Oswestry Disability Index. *Spine (Phila Pa 1976).* 2000;25:2940-
- 42
- 43 19 2953.
- 44
- 45 20 49. EuroQol--a new facility for the measurement of health-related quality of life. *Health Policy.*
- 46
- 47 21 1990;16:199-208.
- 48
- 49
- 50 22 50. Cheung JPY, Luk KDK. The relevance of high-intensity zones in degenerative disc disease. *Int*
- 51
- 52 23 *Orthop.* 2019;43:861-867.
- 53
- 54 24 51. Dudli S, Fields AJ, Samartzis D, Karppinen J, Lotz JC. Pathobiology of Modic changes. *Eur Spine*
- 55
- 56 25 *J.* 2016;25:3723-3734.
- 57
- 58
- 59 26 52. Kyere KA, Than KD, Wang AC, et al. Schmorl's nodes. *Eur Spine J.* 2012;21:2115-2121.
- 60

- 1  
2  
3 1 53. Peters ML, Sommer M, Van Kleef M, Marcus MAE. Predictors of physical and emotional  
4  
5 2 recovery 6 and 12 months after surgery. *Br J Surg*. 2010;97:1518-1527.  
6  
7 3 54. Sullivan M, Bishop S, Pivik J. The pain catastrophizing scale: development and validation.  
8  
9 4 *Psychol Assess*. 1995;7:524-532.  
10  
11 5 55. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand*.  
12  
13 6 Published online 1983. 1983;67:361-370.  
14  
15 7 56. Nicholas MK. The pain self-efficacy questionnaire: Taking pain into account. *Eur J Pain*.  
16  
17 8 2007;11:153-163.  
18  
19 9 57. Reid MC, Williams CS, Gill TM. The relationship between psychological factors and disabling  
20  
21 10 musculoskeletal pain in community-dwelling older persons. *J Am Geriatr Soc*. 2003;51:1092-  
22  
23 11 1098.  
24  
25 12 58. Denison E, Åsenlöf P, Lindberg P. Self-efficacy, fear avoidance, and pain intensity as predictors  
26  
27 13 of disability in subacute and chronic musculoskeletal pain patients in primary health care.  
28  
29 14 *Pain*. 2004;111:245-252.  
30  
31 15 59. Kolahi J, Bang H, Park J. Towards a proposal for assessment of blinding success in clinical trials:  
32  
33 16 Up-to-date review. *Community Dent Oral Epidemiol*. 2009;37:477-484.  
34  
35 17 60. Hróbjartsson A, Emanuelsson F, Thomsen ASS, Hilden J, Brorson S. Bias due to lack of patient  
36  
37 18 blinding in clinical trials. A systematic review of trials randomizing patients to blind and  
38  
39 19 nonblind sub-studies. *Int J Epidemiol*. 2014;43:1272-1283.  
40  
41 20 61. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture  
42  
43 21 (REDCap)-A metadata-driven methodology and workflow process for providing translational  
44  
45 22 research informatics support. *J Biomed Inform*. 2009;42:377-381.  
46  
47 23 62. Manchikanti L, Knezevic NN, Sanapati J, Kaye AD, Sanapati MR, Hirsch J. Is Epidural Injection  
48  
49 24 of Sodium Chloride Solution a True Placebo or an Active Control Agent? A Systematic Review  
50  
51 25 and Meta-Analysis. *Pain Physician*. 2021;24:41-59.  
52  
53  
54  
55  
56  
57  
58  
59 26  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Acknowledgements**

Prof. Roberto S.G.M. Perez had an important role in the initiation of this study and helped substantially with the conception and design of the study. To our greatest regret, he passed away on 07.09.2017.

**Author statement**

EK, PM, JWK, LS, AC and SK designed the study. EK, PM, JWK, JD, LS, AC, PS, DK will conduct the study including patient recruitment and data collection. SK will conduct the data analysis and will conduct the interpretation of the data. JWK and PM drafted the manuscript with important intellectual input from EK, SK, AC, JD, MH, LS, PS, and DK. All authors approved the final manuscript. JWK, PM, JD, and EK will have complete access to the study data.

1  
2  
3 1 **Figure legends**  
4

5 2 **Figure 1**  
6  
7

8 3 1835S GelStix™. From left to right: 18 Gauge Needle, GelStix: dry, after 15 minutes hydration, after  
9  
10 4 30 minutes hydration, after 45 minutes hydration  
11

12 5  
13  
14 6 **Figure 2**  
15

16 7 Study flow chart  
17  
18

19 8  
20

21 9 **Figure 3**  
22

23 10 3A) Using fluoroscopic guidance, the needle is introduced using a standard posterolateral discography  
24  
25 11 approach. 3B) The protective cap is removed from the preloaded implant holder. 3C) The implant  
26  
27 12 holder is threaded onto the proximal end of the introducer needle. 3D ) The holder stylet is pushed  
28  
29 13 so that the implant is driven completely into the introducer needle. 3E) The implant holder is  
30  
31 14 removed. The needle stylet is driven through the needle and bottomed out to deliver the GelStix™  
32  
33 15 completely into the nucleus, keeping the needle tip centered in the nucleus. 3F) The needle tip will  
34  
35 16 keep centered approximately in the nucleus and the procedure will be repeated to insert additional  
36  
37 17 GelStix™.  
38  
39  
40

41 18  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





Figure 1  
1835S GelStix™. From left to right: 18 Gauge Needle, GelStix: dry, after 15 minutes hydration, after 30 minutes hydration, after 45 minutes hydration  
694x517mm (72 x 72 DPI)








Figure 3

3A) Using fluoroscopic guidance, the needle is introduced using a standard posterolateral discography approach. 3B) The protective cap is removed from the preloaded implant holder. 3C) The implant holder is threaded onto the proximal end of the introducer needle. 3D) The holder stylet is pushed so that the implant is driven completely into the introducer needle. 3E) The implant holder is removed. The needle stylet is driven through the needle and bottomed out to deliver the GelStix™ completely into the nucleus, keeping the needle tip centered in the nucleus. 3F) The needle tip will keep centered approximately in the nucleus and the procedure will be repeated to insert additional GelStix™.

338x190mm (96 x 96 DPI)

# Reporting checklist for protocol of a clinical trial.

Based on the SPIRIT guidelines.

## Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the SPIRIT reporting guidelines, and cite them as:

Chan A-W, Tetzlaff JM, Gøtzsche PC, Altman DG, Mann H, Berlin J, Dickersin K, Hróbjartsson A, Schulz KF, Parulekar WR, Krleža-Jerić K, Laupacis A, Moher D. SPIRIT 2013 Explanation and Elaboration: Guidance for protocols of clinical trials. BMJ. 2013;346:e7586

		Page
	Reporting Item	Number
<b>Administrative information</b>		
Title	<a href="#">#1</a> Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	1

1	Trial registration	<a href="#">#2a</a>	Trial identifier and registry name. If not yet registered,	3
2			name of intended registry	
3				
4				
5				
6	Trial registration:	<a href="#">#2b</a>	All items from the World Health Organization Trial	NA
7			Registration Data Set	
8	data set			
9				
10				
11				
12	Protocol version	<a href="#">#3</a>	Date and version identifier	3
13				
14				
15	Funding	<a href="#">#4</a>	Sources and types of financial, material, and other	19
16			support	
17				
18				
19				
20	Roles and	<a href="#">#5a</a>	Names, affiliations, and roles of protocol contributors	1, 27
21				
22	responsibilities:			
23				
24	contributorship			
25				
26				
27				
28	Roles and	<a href="#">#5b</a>	Name and contact information for the trial sponsor	2
29				
30	responsibilities:			
31				
32	sponsor contact			
33				
34	information			
35				
36				
37				
38	Roles and	<a href="#">#5c</a>	Role of study sponsor and funders, if any, in study	2
39			design; collection, management, analysis, and	
40	responsibilities:		interpretation of data; writing of the report; and the	
41			decision to submit the report for publication, including	
42	sponsor and funder		whether they will have ultimate authority over any of	
43			these activities	
44				
45				
46				
47				
48				
49				
50				
51				
52	Roles and	<a href="#">#5d</a>	Composition, roles, and responsibilities of the	NA
53			coordinating centre, steering committee, endpoint	
54	responsibilities:		adjudication committee, data management team, and	
55				
56	committees			
57				
58				
59				
60				

other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)

## Introduction

Background and rationale	<a href="#">#6a</a>	Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention	5-7
Background and rationale: choice of comparators	<a href="#">#6b</a>	Explanation for choice of comparators	5-7
Objectives	<a href="#">#7</a>	Specific objectives or hypotheses	7
Trial design	<a href="#">#8</a>	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, non-inferiority, exploratory)	7
<b>Methods:</b>			
<b>Participants, interventions, and outcomes</b>			
Study setting	<a href="#">#9</a>	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will be collected. Reference to where list of study sites can be obtained	8

1	Eligibility criteria	<a href="#">#10</a>	Inclusion and exclusion criteria for participants. If	8-10
2				
3			applicable, eligibility criteria for study centres and	
4			individuals who will perform the interventions (eg,	
5			surgeons, psychotherapists)	
6				
7				
8				
9				
10				
11	Interventions:	<a href="#">#11a</a>	Interventions for each group with sufficient detail to allow	10-11
12				
13	description		replication, including how and when they will be	
14			administered	
15				
16				
17				
18				
19	Interventions:	<a href="#">#11b</a>	Criteria for discontinuing or modifying allocated	16
20				
21	modifications		interventions for a given trial participant (eg, drug dose	
22			change in response to harms, participant request, or	
23			improving / worsening disease)	
24				
25				
26				
27				
28				
29	Interventions:	<a href="#">#11c</a>	Strategies to improve adherence to intervention	NA
30				
31	adherence		protocols, and any procedures for monitoring adherence	
32			(eg, drug tablet return; laboratory tests)	
33				
34				
35				
36	Interventions:	<a href="#">#11d</a>	Relevant concomitant care and interventions that are	11
37				
38	concomitant care		permitted or prohibited during the trial	
39				
40				
41				
42	Outcomes	<a href="#">#12</a>	Primary, secondary, and other outcomes, including the	12-14
43				
44			specific measurement variable (eg, systolic blood	
45			pressure), analysis metric (eg, change from baseline,	
46			final value, time to event), method of aggregation (eg,	
47			median, proportion), and time point for each outcome.	
48				
49			Explanation of the clinical relevance of chosen efficacy	
50			and harm outcomes is strongly recommended	
51				
52				
53				
54				
55				
56				
57				
58				
59				
60				

Participant timeline	<a href="#">#13</a>	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure)	12, Fig. 2
Sample size	<a href="#">#14</a>	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	14
Recruitment	<a href="#">#15</a>	Strategies for achieving adequate participant enrolment to reach target sample size	8, 17
<b>Methods:</b>			
<b>Assignment of interventions (for controlled trials)</b>			
Allocation: sequence generation	<a href="#">#16a</a>	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	14
Allocation concealment mechanism	<a href="#">#16b</a>	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque,	14-15

1		sealed envelopes), describing any steps to conceal the	
2			
3		sequence until interventions are assigned	
4			
5			
6	Allocation:	<a href="#">#16c</a> Who will generate the allocation sequence, who will	14-15
7			
8	implementation	enrol participants, and who will assign participants to	
9			
10		interventions	
11			
12			
13	Blinding (masking)	<a href="#">#17a</a> Who will be blinded after assignment to interventions	14-15
14			
15		(eg, trial participants, care providers, outcome	
16			
17		assessors, data analysts), and how	
18			
19			
20			
21	Blinding (masking):	<a href="#">#17b</a> If blinded, circumstances under which unblinding is	15
22			
23	emergency	permissible, and procedure for revealing a participant's	
24			
25	unblinding	allocated intervention during the trial	
26			
27			
28			
29	<b>Methods: Data</b>		
30			
31	collection,		
32			
33	management, and		
34			
35	analysis		
36			
37			
38			
39	Data collection plan	<a href="#">#18a</a> Plans for assessment and collection of outcome,	15
40			
41		baseline, and other trial data, including any related	
42			
43		processes to promote data quality (eg, duplicate	
44			
45		measurements, training of assessors) and a description	
46			
47		of study instruments (eg, questionnaires, laboratory	
48			
49		tests) along with their reliability and validity, if known.	
50			
51			
52		Reference to where data collection forms can be found,	
53			
54		if not in the protocol	
55			
56			
57			
58			
59			
60			

1	Data collection plan:	<a href="#">#18b</a>	Plans to promote participant retention and complete	15
2				
3	retention		follow-up, including list of any outcome data to be	
4			collected for participants who discontinue or deviate from	
5			intervention protocols	
6				
7				
8				
9				
10				
11	Data management	<a href="#">#19</a>	Plans for data entry, coding, security, and storage,	15
12			including any related processes to promote data quality	
13			(eg, double data entry; range checks for data values).	
14			Reference to where details of data management	
15			procedures can be found, if not in the protocol	
16				
17				
18	Statistics: outcomes	<a href="#">#20a</a>	Statistical methods for analysing primary and secondary	16-17
19			outcomes. Reference to where other details of the	
20			statistical analysis plan can be found, if not in the	
21			protocol	
22				
23				
24	Statistics: additional	<a href="#">#20b</a>	Methods for any additional analyses (eg, subgroup and	16-17
25	analyses		adjusted analyses)	
26				
27				
28	Statistics: analysis	<a href="#">#20c</a>	Definition of analysis population relating to protocol non-	16
29	population and		adherence (eg, as randomised analysis), and any	
30	missing data		statistical methods to handle missing data (eg, multiple	
31			imputation)	
32				
33				
34	<b>Methods: Monitoring</b>			
35				
36				
37				
38				
39	Data monitoring:	<a href="#">#21a</a>	Composition of data monitoring committee (DMC);	17
40				
41	formal committee		summary of its role and reporting structure; statement of	
42			whether it is independent from the sponsor and	
43				
44				
45				
46				
47				
48				
49				
50				
51				
52				
53				
54				
55				
56				
57				
58				
59				
60				



1		competing interests; and reference to where further	
2		details about its charter can be found, if not in the	
3		protocol. Alternatively, an explanation of why a DMC is	
4		not needed	
5			
6			
7			
8			
9			
10	Data monitoring:	<a href="#">#21b</a> Description of any interim analyses and stopping	17
11	interim analysis	guidelines, including who will have access to these	
12		interim results and make the final decision to terminate	
13		the trial	
14			
15			
16			
17			
18			
19			
20	Harms	<a href="#">#22</a> Plans for collecting, assessing, reporting, and managing	14
21		solicited and spontaneously reported adverse events	
22		and other unintended effects of trial interventions or trial	
23		conduct	
24			
25			
26			
27			
28			
29			
30	Auditing	<a href="#">#23</a> Frequency and procedures for auditing trial conduct, if	NA
31		any, and whether the process will be independent from	
32		investigators and the sponsor	
33			
34			
35			
36			
37			
38	Ethics and		
39	dissemination		
40			
41			
42			
43	Research ethics	<a href="#">#24</a> Plans for seeking research ethics committee /	19
44	approval	institutional review board (REC / IRB) approval	
45			
46			
47			
48	Protocol	<a href="#">#25</a> Plans for communicating important protocol	19
49	amendments	modifications (eg, changes to eligibility criteria,	
50		outcomes, analyses) to relevant parties (eg,	
51		investigators, REC / IRBs, trial participants, trial	
52		registries, journals, regulators)	
53			
54			
55			
56			
57			
58			
59			
60			

Consent or assent	<a href="#">#26a</a>	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	19
Consent or assent: ancillary studies	<a href="#">#26b</a>	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	NA
Confidentiality	<a href="#">#27</a>	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	19
Declaration of interests	<a href="#">#28</a>	Financial and other competing interests for principal investigators for the overall trial and each study site	19
Data access	<a href="#">#29</a>	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	27
Ancillary and post trial care	<a href="#">#30</a>	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	NA
Dissemination policy: trial results	<a href="#">#31a</a>	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	15-16

1	Dissemination policy: <a href="#">#31b</a>	Authorship eligibility guidelines and any intended use of	NA
2			
3	authorship	professional writers	
4			
5			
6	Dissemination policy: <a href="#">#31c</a>	Plans, if any, for granting public access to the full	NA
7			
8	reproducible	protocol, participant-level dataset, and statistical code	
9			
10	research		
11			
12			
13			

14 **Appendices**

15			
16			
17	Informed consent	<a href="#">#32</a> Model consent form and other related documentation	appendix
18			
19	materials	given to participants and authorised surrogates	1
20			
21			
22	Biological specimens	<a href="#">#33</a> Plans for collection, laboratory evaluation, and storage of	NA
23			
24		biological specimens for genetic or molecular analysis in	
25			
26		the current trial and for future use in ancillary studies, if	
27			
28		applicable	
29			
30			
31			

32 The SPIRIT Explanation and Elaboration paper is distributed under the terms of the Creative  
33 Commons Attribution License CC-BY-NC. This checklist was completed on 20. May 2021 using  
34 <https://www.goodreports.org/>, a tool made by the [EQUATOR Network](#) in collaboration with  
35 [Penelope.ai](#)  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

# BMJ Open

## **Efficacy of the Gelstix nucleus augmentation device for the treatment of chronic discogenic low back pain: protocol for a randomised, sham-controlled, double-blind, multicentre trial**

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-053772.R1
Article Type:	Protocol
Date Submitted by the Author:	11-Feb-2022
Complete List of Authors:	Koetsier, Eva; Neurocenter of Southern Switzerland, EOC, Pain Management Center; Università della Svizzera Italiana, Faculty of Biomedical Sciences van Kuijk, Sander; Maastricht Universitair Medisch Centrum+, Clinical Epidemiology and Medical Technology Assessment Maino, P.; Neurocenter of Southern Switzerland, EOC, Pain Management Center; Università della Svizzera Italiana, Faculty of Biomedical Sciences Dukanac, J.; Neurocenter of Southern Switzerland, EOC, Pain Management Center Scascighini, L.; University of Applied Sciences and Arts of Southern Switzerland, Department of Health Sciences Cianfoni, A.; Neurocenter of Southern Switzerland, EOC, Service of Diagnostic and Interventional Neuroradiology; Inselspital University Hospital Bern, Dept. of Neuroradiology Scarone, P.; Neurocenter of Southern Switzerland, EOC, Clinic of Neurosurgery Kuhlen, D.E.; Neurocenter of Southern Switzerland, EOC, Clinic of Neurosurgery Hollman, M; Amsterdam UMC Locatie AMC, Department of Anesthesiology Kallewaard, Jan-Willem; Rijnstate, Anesthesiology
<b>Primary Subject Heading</b>:	Anaesthesia
Secondary Subject Heading:	Radiology and imaging, Rheumatology
Keywords:	Back pain < ORTHOPAEDIC & TRAUMA SURGERY, PAIN MANAGEMENT, Musculoskeletal disorders < ORTHOPAEDIC & TRAUMA SURGERY

SCHOLARONE™  
Manuscripts

1

2

31Efficacy of the Gelstix nucleus augmentation device for the treatment of chronic discogenic low back

4

52

6pain: protocol for a randomised, sham-controlled, double-blind, multicentre trial

7

83

9

104E. Koetsier<sup>1,2</sup>, S.M.J. van Kuijk<sup>3</sup>, P. Maino<sup>1,2</sup>, J. Dukanac<sup>1</sup>, L. Scascighini<sup>4</sup>, A. Cianfoni<sup>5,6</sup>, P. Scarone<sup>7</sup>, D.E.

11

125

13Kuhlen<sup>7</sup>, M.W. Hollmann<sup>8</sup>, J.W. Kallewaard.<sup>8,9</sup>

146

15

167

17

188

191. Pain Management Center, Neurocenter of Southern Switzerland, EOC, Lugano, Switzerland

20

219

222. Faculty of Biomedical Sciences, Università della Svizzera Italiana, Lugano, Switzerland

23

2410

253. Department of Clinical Epidemiology and Medical Technology Assessment, Maastricht University

26

2711

28Medical Centre+, Maastricht, the Netherlands

29

3012

314. Department of Health Sciences, University of Applied Sciences and Arts of Southern Switzerland,

32

3313

34Manno, Switzerland

35

3614

375. Service of Diagnostic and Interventional Neuroradiology, Neurocenter of Southern Switzerland,

38

3915

40EOC, Lugano, Switzerland

41

4216

436. Dept. of Neuroradiology, Inselspital University Hospital Bern, Switzerland

44

4517

467. Clinic of Neurosurgery, Neurocenter of Southern Switzerland, EOC, Lugano, Switzerland

47

4818

498. Department of Anesthesiology, Amsterdam UMC, location Academic Medical Center, Amsterdam,

50

5119

52Netherlands

53

5420

559. Department of Anesthesiology and Pain Management Arnhem, Rijnstate Hospital, Velp, the

56

5721

58Netherlands

59

6022

23Word count: 4115

24Number of tables: 0

25Number of figures: 3

26

1 Address correspondence and reprint requests to:

2 Dr. Eva Koetsier, Pain Management Center, Neurocenter of Southern Switzerland, EOC, Lugano,  
3 Switzerland.

4 Ospedale Regionale di Lugano, Sede Ospedale Italiano, via Pietro Capelli 1, 6962, business telephone  
5 number: +41 (0)91 8117590, E-mail: [evakoetsier6@gmail.com](mailto:evakoetsier6@gmail.com), [eva.koetsier@eoc.ch](mailto:eva.koetsier@eoc.ch).

6  
7 Sponsor contact information:

8 Pain Management Center, Neurocenter of Southern Switzerland, EOC, Lugano, Switzerland. Ospedale  
9 Regionale di Lugano, Sede Ospedale Italiano, via Pietro Capelli 1, 6962, Lugano, Switzerland, business  
10 telephone number: +41 (0)91 8117590, E-mail: [eva.koetsier@eoc.ch](mailto:eva.koetsier@eoc.ch).

1

2

31 ABSTRACT

4

52 Introduction

6

73 Discogenic pain is the cause of pain in 26-40% of patients with for low back pain. Consensus about

8

94 treatment of chronic discogenic low back pain is lacking and most treatment alternatives are

10

115 supported by limited evidence. The percutaneous implantation of hydrogels into the nucleus

12

136 pulposus represents a promising regenerative intradiscal therapy. The hydrogel ‘GelStix™’ is

14

157 composed primarily of hydrolyzed polyacrylonitrile and acts as a reservoir of hydration, producing

16

178 increased pressure and improved pH balance, potentially leading to disc preservation. We

18

199 hypothesize that treatment with GelStix™ will lead to greater reduction in pain intensity at six

20

2110 months post-treatment compared to patients receiving sham treatment.

22

2311 Methods and analysis

24

2512 This is a parallel group, randomized sham-controlled double-blind, multicentre trial to assess whether

26

2713 the GelStix™ device is superior to sham in reducing pain intensity in patients with chronic discogenic

28

2914 low back pain. The study will be conducted in two regional hospitals in Europe. Seventy-two

30

3115 participants will be randomized in a 1:1 ratio. The primary outcome will be the change in pain intensity

32

3316 between preoperative baseline and at six months post-intervention. Secondary outcomes were

34

3517 disability, quality of life, the patient’s global impression of change scale, the use of pain medication,

36

3718 and the disc degeneration process assessed by means of MRI. For change in pain intensity, disability,

38

3919 health related quality of life, and disc height, mean values will be compared between groups using

40

4120 linear regression analysis, adjusted for treatment centre.

42

4321 Ethics and dissemination

44

4522 Ethics approval was obtained from the Ethics Committee of the Canton Ticino, Switzerland (CE2982)

46

4723 and by the Medical Ethical Committee Arnhem-Nijmegen, the Netherlands (2016-2944). All patients

48

4924 that agree to participate will be asked to sign an informed consent form. Results will be disseminated

50

5125 through international publications in peer reviewed journals, in addition to international conference

52

5326 presentations.

54

55

56

57

58

59

60

1

2 **Trial registration number** NCT027639563 **Protocol version** 7.1, 18/11/20204 **Keywords** Back pain, pain management, musculoskeletal disorders

5

6

7 **ARTICLE SUMMARY**8 **Strengths and limitations of this study**

9 ► This will be the first prospective, randomized, controlled, multicentre trial assessing effectivity and  
10 safety of the GelStix™ Nucleus Augmentation Device compared to a sham control in patients with  
11 lumbar discogenic pain that had no benefit from conservative care.

12 ► Means to reduce risk of bias are implemented, which includes an a-priori sample size calculation,  
13 an explicitly stated primary hypothesis to be tested, methodological rigor, double-blinding,  
14 randomization, adequate concealment of group allocation and the assessment of the success of  
15 blinding in participants and observers.

16 ► This is also the first study that assesses the disc degeneration process and disc height by means of  
17 Magnetic Resonance Imaging (MRI) one year after GelStix™ implantation versus sham.

18 ► All participants will also be treated according to a protocolized physiotherapy.

19 ► The limitations are those inherent to a prospective, randomized sham-controlled double-blind  
20 study, including strict exclusion criteria and thus limited generalizability (e.g., protrusions in contact  
21 with any nerve root at the symptomatic level or >5mm, an insufficient number of patients, and  
22 adherence to a strict protocol that does not necessarily reflect real word daily practice).



1

2

31INTRODUCTION

4

52Background and rationale

6

7

83Discogenic low back pain is characterized by persistent, predominantly centralized axial low back

9

104pain that worsens with axial loading. It is associated with intervertebral disc degeneration without

11

125herniation,<sup>1–4</sup> and is thought to be the cause of pain in 26–40% of patients consulting a physician for

13

146low back pain.<sup>5–9</sup> The water-binding capabilities of the intervertebral disc diminish with aging<sup>10</sup>

15

167leading to progressive shrinking of the nucleus pulposus and loss of elasticity.<sup>10–13</sup> The cartilaginous

17

188endplate vascular flow decreases due to a progressive loss in vascularization leading to accumulation

19

209of cellular waste products, and an increasingly acidic environment.<sup>10,14</sup> A low pH around the discus is

21

2210associated with discogenic pain.<sup>15,16</sup>

23

2411Medical history, physical examination, and imaging (e.g. magnetic resonance imaging (MRI)) provide

25

2612inadequate sensitivity and specificity to accurately diagnose discogenic pain.<sup>17–21</sup> Despite an ongoing

27

2813debate, moderate evidence supports diagnostic accuracy of provocative discography.<sup>19,22–24</sup> While

29

3014previous studies suggest that high-pressure provocative discography may accelerate disc

31

3215degeneration,<sup>25–27</sup> a recently published study suggests that low-pressure provocative discography,

33

3416performed according to International Association for the Study of Pain (IASP) criteria, does not

35

3617accelerate disc degeneration.<sup>28</sup>

37

3818Consensus about treatment of chronic discogenic low back pain is lacking and the majority of

39

4019treatment alternatives is supported by limited evidence.<sup>1,4</sup> Conservative management includes anti-

41

4220inflammatory drugs, physiotherapy, and multidisciplinary biopsychosocial rehabilitation.<sup>29</sup> If

43

4421conservative treatment fails, (minimally) invasive treatments are considered.<sup>1</sup> Most minimally

45

4622invasive treatments, such as intradiscal injections (e.g. with methylene blue) and thermal

47

4823intradiscal/annular techniques (intradiscal electrothermal therapy (IDET), have been abandoned

49

5024because of poor evidence.<sup>30–32</sup> A recent systematic review concluded that most minimal invasive

51

5225treatments for discogenic low back have very low evidence; only biacuplasty has moderate evidence

53

5426for a subgroup of patients with discogenic low back pain.<sup>33</sup>

55

56

57

58

59

60

1 Fusion surgery and total disc replacement, although contemplated as possible therapies in some  
2 cases, are invasive interventions associated with risk of adjacent segment disorder and morbidity.<sup>4,34</sup>  
3 In addition, fusion surgery is not superior to conservative treatment with multidisciplinary  
4 biopsychosocial rehabilitation and physiotherapy.<sup>35,36</sup> Recently, with the emergence of new  
5 frequencies (burst, dorsal root ganglion stimulation, high frequency-10Hz), low back pain has  
6 become a good treatment option for neuromodulation. Considering the fact that neuromodulation is  
7 a more invasive treatment the need is great to find evidence for minimal invasive treatment for  
8 chronic discogenic low back pain.<sup>37,38</sup>  
9 Therefore, treatment options filling the gap between conservative care and invasive surgical  
10 intervention are urgently needed. Currently the first studies are published showing effect of the use  
11 of platelet-rich plasma (PRP) and mesenchymal signaling cells (MSCs) for discogenic pain. Notably, no  
12 intervention has multiple RCT's published yet.<sup>39</sup> The implantation of hydrogels into the nucleus  
13 pulposus represents a promising regenerative intradiscal therapy, in particular in patients with early  
14 or moderate disc degeneration not responding to conservative care.<sup>40,41</sup> The hydrogel containing  
15 'GelStix™ Nucleus Augmentation Device' (hereafter called GelStix™) is composed primarily of  
16 hydrolyzed polyacrylonitrile (HPAN). The GelStix™ is shaped in the form of an elongated matchstick  
17 and can be inserted percutaneously into the nucleus through a needle. Once implanted, the GelStix™  
18 absorbs the body's own fluids and expands around tenfold in volume (see Fig. 1).

19  
20 **Insert here Figure 1**

21  
22 The GelStix™ material acts as a reservoir of permanent hydration of the intervertebral disc,  
23 producing increased pressure, and improved fluid exchange and pH balance, leading to disc  
24 preservation.<sup>42</sup> Results of previous non-controlled studies suggest that GelStix™ implantation leads to  
25 a significant pain and disability relief four weeks after implantation in patients with discogenic  
26 pain.<sup>43,44</sup>

## 2 Objectives

The primary outcome will be the change in pain intensity between preoperative baseline and at six months post-intervention. Secondary outcomes include disability, quality of life outcome measures, the patient's global impression of change (PGIC) scale, the use of pain medication, and the disc degeneration process assessed by means of MRI.

9 We hypothesize that treatment with GelStix™ will lead to greater reduction in pain intensity at six  
10 months post-treatment compared to patients receiving sham treatment.

## 2 Trial design

3 This is a parallel group, randomized sham-controlled double-blind, multicentre trial to assess whether  
4 the GelStix<sup>tm</sup> device is superior to sham in reducing pain intensity in patients with chronic discogenic  
5 low back pain. Patients are randomly allocated in a 1:1 ratio. Figure 2 provides a flow diagram of the  
6 progress through the enrolment, intervention allocation, follow-up, and data analysis phases of the  
7 trial.

9      **Insert here Figure 2**

## METHODS AND ANALYSIS

This protocol has been written in accordance with the Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT) checklist. The study will be conducted in two regional hospitals in Europe: the Pain Management Center, Neurocenter of Southern Switzerland, Lugano, Switzerland, and the Department of Anaesthesiology and Pain Management Arnhem, Rijnstate Hospital, Arnhem, the Netherlands. Recruitment started in April 2016 and we included 42 participants till now. We expect to complete the study in 2025.

### Participants

The target population is represented by patients suffering from discogenic low back pain with a baseline numeric rating scale (NRS) pain score  $\geq 5/10$  following at least twelve weeks conservative care.

#### Inclusion criteria:

- 18-66 years of age
- Lumbar DDD on MRI scan with Pfirrmann grade<sup>45</sup> 2, 3 or 4
- Discogenic pain confirmed by positive discography\* of one or maximum two lumbar disc levels, and one negative control level
- Persistent predominant, nociceptive low back pain with a NRS score of  $\geq 5/10$ , that worsens with axial loading and improves with recumbence of at least 12 weeks duration
- Failure to have symptoms resolved or reduced following at least 12 weeks conservative care (drug therapy and/or physiotherapy)
- Negative medial branches block results
- Legally competent and able to understand the nature, scope and aim of the clinical investigation

1		
2		
3	1	
4		
5	2	Exclusion criteria:
6		
7		
8	3	• Radiculopathy caused by nerve root compression
9		
10	4	• Frank herniations, extruded or sequestered fragments, bulge/protrusions in contact with any
11		
12	5	nerve root at the symptomatic level or >5mm in antero-posterior dimension
13		
14	6	• Greater than grade 4 annular tear (Adams scale) <sup>46</sup>
15		
16		
17	7	• Disc height less than 3mm at the symptomatic level
18		
19	8	• Severe symptomatic central, foraminal or lateral recess stenosis, spondylolysis,
20		
21	9	spondylolisthesis greater than I out of IV, acute fractures, or ankylosing spondylitis at any
22		
23	10	lumbar disc level
24		
25		
26	11	• Coagulopathy or oral anticoagulant therapy (except low-dose acetylsalicylic acid) in
27		
28	12	conditions that do not allow for a temporary discontinuation
29		
30	13	• Active infection, systemically or localized
31		
32		
33	14	• Any disease process or condition that may make the effect of the treatment difficult to
34		
35	15	evaluate (e.g. cancer, substance abuse, etc.)
36		
37	16	• Previous surgery at any lumbar disc level
38		
39		
40	17	• Body Mass Index (BMI) of ≥ 35 kg/m <sup>2</sup>
41		
42	18	• Females of childbearing age that are known to be pregnant or wishing to be pregnant during
43		
44	19	the study
45		
46	20	• Psychological disorders or factors that may impact upon treatment outcomes or compliance
47		
48	21	(e.g. severe depressions)
49		
50		
51	22	• Participation in any other interventional study at the same time
52		
53	23	
54		
55	24	*Procedure of provocative discography
56		
57		
58	25	Provocative discography will be performed by an experienced pain physician under strict sterile
59		
60	26	conditions. Thirty minutes before the intervention, intravenous antibiotics for prophylaxis will be

administered. The patient will be positioned in the prone position on an X-ray permeable table. After subcutaneous anaesthetic injection of 2 ml mg of lidocaine 1%, the nucleus will be accessed with the two-needle technique with a 25-27 Gauge needle through the transforaminal, posterolateral approach, according to the technique described by Kallewaard et al.<sup>3</sup> Fluoroscopy will be used to identify spinal levels, guide the needle, and to confirm final needle position. The following variables will be monitored during the injection of the contrast solution: the opening pressure (the pressure at which contrast is first visible in the disc), the provocation pressure (the pressure greater than the opening pressure at which complaints of pain arise), and the peak pressure or the final pressure at the end of the procedure. Additionally, the total volume of the injected contrast solution, the Adams scale,<sup>46</sup> and the pain score measured by NRS per disc level will be recorded.

The procedure, per level, is continued until:<sup>3</sup>

- Concordant pain is reproduced at a level of  $\geq 7/10$  and/or
- The volume infused reaches 3.0 mL and/or
- The pressure rises to 50 psi above opening pressure

According to the guidelines of the IASP,<sup>47</sup> the symptomatic level and the one adjacent level are examined. A disc is only considered to be positive if concordant pain can be induced at the target level (symptomatic level); with an intensity of this pain of at least NRS 7, reproduced by a pressure of less than 50 psi above opening pressure; and if the control level is negative for provocation of pain. A control disc is considered a critical element for defining a positive discography, as it serves as an internal patient control disc and as a possible indicator of central sensitization.

## Interventions

### The GelStix™ implantation

For each participant, up to two levels will be treated. The CE marked GelStix™ Nucleus Augmentation Device system (STX-1835S, Replication Medical, Inc. – Cranbury, NJ, USA), will be implanted by an

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 experienced pain physician familiar with the transforaminal posterolateral discography approach  
2 described above. The GelStix™ insertion will be performed under local anaesthesia with a single  
3 needle technique through the procedure-specific 18 Gauge needle (18GTXX165mm, Replication  
4 Medical, Inc. – Cranbury, NJ, USA). Up to three GelStixs will be implanted at each symptomatic disc  
5 level. Once the needle tip is located in the centre of the nucleus, the stylet will be removed from the  
6 needle. Then, the protective cap is removed from the preloaded GelStix™ holder and the GelStix™  
7 holder is threaded onto the proximal end of the introducer needle. The holder stylet is pushed,  
8 driving the GelStix™ completely into the introducer needle. The implant holder will then be removed  
9 and the needle stylet ('blunt push rod needle') is driven through the needle and bottomed out to  
10 deliver the GelStix™ completely into the nucleus, keeping the needle tip centred in the nucleus (fig.  
11 3A-3F). The procedure will be repeated to insert additional GelStix™. When resistance rises adding a  
12 second or third GelStix™, further insertion is discontinued. At the end of the procedure, the needle  
13 will be withdrawn, and a sterile bandage will be applied to the insertion site.

15 **Insert here Figure 3**

17 The sham intervention

18 For the sham intervention the symptomatic discs will be injected with 1 ml of saline (NaCl 0.9%).  
19 Intradiscal saline injection (1 mL NaCl 0.9% ) is safe<sup>48</sup> and has been used as a control/sham  
20 intervention in other randomized controlled <sup>30,49,50</sup>

22 Concomitant treatment

23 Starting two weeks after the intervention, participants of both study groups will be prescribed  
24 physiotherapy according to a study specific protocol. Session frequency will be once a week, for nine  
25 weeks. An experienced musculoskeletal physiotherapist will assess the patient before starting the  
26 post-intervention protocol, in order to determine the starting level for the exercises. Motor control

1 and stabilization exercises will be instructed to the patients and they will get a leaflet with pictures of  
2 the exercises to perform at home/at work. Individual exercises include training of the deep  
3 abdominal muscles with the lumbar multifidus and the transversus abdominis. Moreover, to restore  
4 the function of the core muscles, all directions and their muscular chains will be trained. All patients  
5 will be instructed as to how to do exercises at home and will be asked to continue these exercises  
6 three times a week for six months. Continuation or modification of pain medication is permitted  
7 during the study period of twelve months.

### 9 **Outcome measures**

10 The primary outcome is the change in pain intensity, assessed by means of a pain diary, between  
11 preoperative baseline and at six months post-intervention in the GelStix<sup>tm</sup>-treated compared to the  
12 sham-treated group. Pain intensity will be assessed employing an 11-point (i.e. 0–10) NRS with 0  
13 meaning ‘no pain’ and ‘10’ meaning ‘worst possible pain’.<sup>51</sup> Three times daily pain scores will be  
14 assessed for five consecutive days around the intended measurement time. The mean NRS scores on  
15 the pain diary will furthermore be measured at one week, and one, three, and twelve months.

17 The secondary outcomes include:

- 18 - Disability, using the Oswestry Disability Index (ODI). The ODI is completed at baseline, and at  
19 three, six and twelve months. The ODI is a self-administered questionnaire, assessing the  
20 patient’s level of pain and function during basic activities of daily living such as walking,  
21 personal care, standing, sleeping, etc.<sup>52</sup>
- 22 - Quality of life (QoL), quantified with the European Quality of Life Five Dimension Five Level  
23 Scale (EQ-5D-5L). The EQ-5D-5L will be completed at baseline and at three, six and twelve  
24 months. This questionnaire assesses health related quality of life in terms of five dimensions:  
25 mobility, self-care, usual activities, pain/discomfort, and anxiety/depression.<sup>53</sup> Additionally,  
26 the EuroQol Visual Analogue Scale (EQ VAS) records the respondent’s self-rated health on a 20



1  
2  
3 1 cm vertical, visual analogue scale with endpoints labelled ‘the best health you can imagine’  
4  
5 2 and ‘the worst health you can imagine’.  
6  
7 3 - The Patient’s Global Impression of Change (PGIC) scale will be measured at three, six and  
8  
9 twelve months. This scale assesses the patient's own evaluation of improvement or  
10 4  
11 deterioration over time on a 7-point Likert Scale rated from ‘very much improved’ to ‘very  
12 5  
13 much worse’.  
14 6  
15  
16 7 - The use of pain medication will be assessed as the intake of analgesics at baseline, at one  
17 8  
18 week, and at one, three, six and twelve months.  
19 9  
20 - The disc degeneration process will be assessed by means of MRI twelve months after  
21 10  
22 treatment compared to baseline. Pfirrmann grade,<sup>45</sup> disc height, and the presence of high  
23 11  
24 intensity zones (HIZ),<sup>54</sup> Modic signs,<sup>55</sup> and Schmorl’s nodes<sup>56</sup> will be recorded.  
25 12  
26  
27  
28  
29  
30 13 Additionally, to assess the association between pain catastrophizing, surgical fear, state of  
31  
32 14 depression and long-term outcome the following additional patient-reported outcome measures  
33  
34 15 (PROMs) will be registered at baseline. Pain catastrophizing, defined as an exaggerated negative  
35  
36 16 interpretation of the meaning of pain, will be measured by the Pain Catastrophizing Scale (PCS).  
37  
38 17 Higher pain catastrophizing before intervention are related to lower perceived recovery.<sup>57,58</sup> Surgical  
39  
40 18 fear will be measured by the Surgical Fear Questionnaire (SFQ) as a predictor of physical and  
41  
42 19 emotional recovery.<sup>57</sup> State of depression will be assessed by the Hospital Anxiety and Depression  
43  
44 20 Scale (HADS), a self-administered questionnaire developed to detect states of anxiety and depression  
45  
46 21 in hospital out-patient clinics.<sup>59</sup> Moreover, pain self-efficacy will be assessed employing the Pain Self-  
47  
48 22 Efficacy Questionnaire-I (PSEQ-I). This patient self-reported measurement instrument evaluates pain  
49  
50 23 self-efficacy beliefs,<sup>60</sup> i.e. the degree of confidence a patient has in performing regular daily activities  
51  
52 24 despite of pain. The presence of low levels of pain self-efficacy has been shown to be associated with  
53  
54 25 high levels of disability in patients experiencing pain.<sup>61,62</sup>  
55  
56  
57  
58  
59 26  
60

The following additional data will be collected at baseline: sex, age, weight, height, smoking habits, previous treatment of discogenic pain, and neurological examination. Employment status baseline and at six and twelve months will be recorded. The proportion of patients unable to return to work will be an additional measure of efficacy of the treatment.

The success of blinding will be assessed at the end of the trial. Before unblinding, the patients and the blind observers will be asked to guess the patients' treatment and the answers will be compared with the actual treatments administered. Successful blinding procedures can reduce bias in clinical trials.<sup>63,64</sup>

The safety outcome of this study is the incidence and severity of complications and adverse events (AE's) including procedure-related complications at any time point in the study. The main expected adverse device effects are infection (local or discitis), bleeding, nerve damage and/or limited motion as a result of the procedure.

### Sample size

Twenty-eight patients per group will be required to have 80% power to detect a minimally clinically relevant difference of 1.5 points on the NRS between groups, with an estimated standard deviation (SD) of 2, based on the pooled SD of NRS scores of similar patients in the RCT of Kallewaard et al.,<sup>30</sup> and testing with an alpha of 5% (two-tailed). With an expected drop-out rate of about 20%, a total of 72 patients will be randomized.

### Randomization

The Project Manager of the Clinical Trial Unit of the Ente Ospedaliero Cantonale (CTU-EOC), Bellinzona, Switzerland, will be in charge for computer generated block randomization lists stratified

by centre (blocks of 4). The Project Manager will act as an independent person, not involved in any other aspect of the trial except administrative/financial issues. The study is patient- and observer-blinded, while the physician performing the study intervention will necessarily be aware of the treatment allocation. A web-based access to patient allocation codes will be provided to the physician in charge for GelStix<sup>tm</sup>/placebo injection. The treating team will be instructed not to communicate allocation to GelStix<sup>tm</sup> or placebo in any way, both to the patient and to other trial personnel. The “assessors”, i.e., the investigators in charge for efficacy and safety assessments and the research nurses that may be in charge for questionnaires collection, and the personnel in charge of monitoring/data review and analysis will have no access to the randomization lists and will receive no information about patient treatment for the entire duration of the study. For patients still experiencing substantial discogenic pain at six months, the code can be broken at their request (after the assessment of the success of blinding). The patients initially allocated to the control group are then given the opportunity to cross-over to the GelStix<sup>tm</sup> treatment. Any other code breaks should occur only in circumstances when knowledge of the actual treatment is absolutely essential for further management of the patient e.g., in case of important AE’s to ensure the most appropriate patient management.

**Data collection and management**

Study data will be collected on a case report form by the research team and will be entered in a research electronic data capture (REDCap) database.<sup>65</sup> The data will be associated to a unique trial identification number per patient. The database will be double-checked for missing data and data entry errors. The data from the REDCap database will be imported automatically in the latest version of R, a language for statistical computing. All study data will be archived for at least of 15 years after study termination.

1

## 2 Statistical methods

3 Baseline characteristics will be described stratified by treatment allocation as mean and standard  
4 deviation or median and first and third quartile, and as count and percentage, as appropriate. In case  
5 of over 5% of missing data, we will use multiple imputation with fully conditional specification to  
6 impute the dataset. The number of imputations will be set to the percentage of incomplete patients.  
7 All subsequent analyses will be performed according to the intention to treat principle. A “per  
8 protocol” analysis will also be performed, excluding patients who are not evaluable for the primary  
9 endpoint because of dropout (e.g., consent withdrawal before completion of the six months  
10 observation period). Frequency and type of AE’s and complications during the study will be described  
11 in the final report. Dropouts will be replaced up to the number of evaluable patients defined in the  
12 sample size calculation.

13 The primary outcome is change in pain (NRS) at six months compared to baseline. Mean values will  
14 be compared between groups using linear regression analysis, adjusted for treatment centre. In case  
15 of imbalance of baseline characteristics as judged by the trial steering committee, regression  
16 analyses will be further adjusted for potential confounders. This adjustment will be performed as  
17 stratified randomization induces correlated observations, which should be accounted for. By  
18 adjusting for treatment center, the analyses yield correct p-values and confidence intervals with the  
19 correct coverage, and results in more power compared to unadjusted analyses.<sup>66</sup>

20 Change from baseline in pain at other follow-up moments and change from baseline in continuous  
21 secondary outcome measures (i.e., disability (ODI) and health related quality of life [EQ-5D-5L], and  
22 disc height) will be analysed in a similar manner. PGIC scores will be dichotomized by taking “very  
23 much improved” and “much improved” to indicate treatment success. Pfirrmann grade will be  
24 dichotomized into grade 1 or 2 versus more than 2. Success rates on the PGIC, dichotomized  
25 Pfirrmann grade, and the presence of HIZ, Modic changes, and Schmorl’s nodes will be compared  
26 between groups using logistic regression analysis adjusted for center.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 Univariable and multivariable logistic regression will be used to quantify crude and adjusted  
2 associations between PCS, SFQ, HADS, and PSEQ-I and treatment success. These analyses will be  
3 considered exploratory. The success of blinding will be assessed using the Sign test, testing whether  
4 the percentage of correct guesses differs from that expected by chance (i.e. 50%).  
5

6 **Monitoring**

7 The research project will be monitored by a certified clinical monitor, which will review the data  
8 quality and will ensure that study activities are carried out in accordance with the protocol, good  
9 clinical practice and applicable regulatory requirements. This being a novel treatment method, a  
10 blinded interim analysis for futility will be planned for the primary outcome measure at T3 months  
11 after 40 patients (i.e. 20 in each arm of the study) have been enrolled. The study will be terminated  
12 in case the experimental arm performs significantly worse (as based on independent samples t-test  
13 or Mann-Whitney-U test) *and* the difference between groups is clinically relevant (i.e. 2 points or  
14 more on the NRS).  
15  
16

17 **Limitations of the study**

18 The limitations are those inherent to a prospective, randomized, sham-controlled study, including  
19 difficulty in recruiting patients due to potential patient refusal and strict exclusion criteria (e.g.,  
20 protrusions in contact with any nerve root on the symptomatic level or >5mm), an insufficient  
21 number of patients, and adherence to a strict protocol that does not necessarily reflect real world  
22 daily practice. Recently performed strategies for achieving adequate participant enrolment to reach  
23 target sample size are the drafting and dispersal of an informative letter to referral colleagues in  
24 Switzerland and in the Netherlands, the introduction of a back pain treatment algorithm in the Pain  
25 Management Center in Lugano, indicating a clear algorithm to follow after negative medial branch  
26 block tests, indicating also the possibility for inclusion in the GelStix™ study.

Another limitation of this trial is the question whether intradiscal saline injection is a true placebo, as it may have active effects. For example, a recently published systematic review and meta-analysis of Manchikanti et al. showed that epidurally administered saline and saline with steroids may be both effective in managing low back and lower extremity pain.<sup>67</sup> On the other hand, saline has been routinely used as a sham intervention in several other intradiscal treatment studies such as the randomized controlled trial (RCT) of Kallewaard et al.,<sup>30</sup> which compared intradiscal methylene blue plus lidocaine to intradiscal saline plus lidocaine injection, and two the RCT's of Cao et al.<sup>49</sup> and Khot et al.<sup>50</sup> comparing intradiscal corticosteroid to saline injection in the treatment of discogenic low back pain. To reduce the risk of a bias due to the uncertainty saline injection being a true placebo, a third 'no treatment group' (receiving only physiotherapy treatment) could be added to this study. However, we regard adding a third 'no treatment group' to this study not feasible, mainly because of the expected difficulties in patient recruitment.

#### **Patient and public involvement**

Patient with discogenic pain were involved at several stages of the trial, including the design and conduct of the trial. We carefully assessed the burden of the trial interventions on these patients. We will disseminate the main results to trial participants and will seek patient and public involvement in the development of an appropriate method of dissemination.

1  
2  
3 1 **ETHICS AND DISSEMINATION**  
4

5 2  
6  
7 3 **Research ethics approval and consent to participate**  
8  
9  
10 4 This trial has been approved by the Research Ethics Committee of the Canton Ticino, Switzerland (CE  
11  
12 5 2982) and by the Medical Ethical Committee Arnhem-Nijmegen, the Netherlands (2016-2944). All  
13  
14 6 patients that agree to participate will sign an informed consent form provided by the independent  
15  
16 7 observer. Any amendment to the protocol must as well be approved by this institution.  
17  
18  
19 8

20  
21 9 **Confidentiality**  
22  
23 10 Individual subject medical information obtained as a result of this study is considered confidential  
24  
25 11 and disclosure to third parties is prohibited. Subject confidentiality will be further ensured by utilizing  
26  
27 12 subject identification code numbers. Direct access to source documents will be permitted for  
28  
29 13 purposes of data review by authorized personnel involved in the trial and inspections. Patients’  
30  
31 14 identity will not be disclosed to the person in charge for the statistical analysis and will not appear in  
32  
33 15 any publication or public presentation of the study results.  
34  
35  
36  
37 16

38  
39 17 **Dissemination**  
40  
41 18 Results will be disseminated through international publications in peer reviewed journals, in addition  
42  
43 19 to international conference presentations.  
44  
45  
46 20

47  
48 21 **Funding statement**  
49  
50 22 This research is not supported by a topic-specific grant from any funding agency in the public,  
51  
52 23 commercial or not-for-profit sectors. Research personnel will in part be paid by previous grants that  
53  
54 24 were not awarded for this specific study. The manufacturing company Replication Medical, Inc. will  
55  
56 25 cover the costs of the GelStix™ material. This will be an unrestricted support and the manufacturer  
57  
58  
59  
60

1 will not be involved in study design, data collection, analysis, interpretation, and  
2  
3  
4  
5 2 reporting/publication.  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

3

4 **Competing interest**

5 None declared

6

For peer review only



References

1. Zhao L, Manchikanti L, Kaye AD, Abd-Elseyed A. Treatment of Discogenic Low Back Pain: Current Treatment Strategies and Future Options—a Literature Review. *Curr Pain Headache Rep.* 2019;23:86.

2. Peng BG. Pathophysiology, diagnosis, and treatment of discogenic low back pain. *World J Orthop.* 2013;4:42-52.

3. Kallewaard JW, Terheggen MAMB, Groen GJ, et al. 15. Discogenic Low Back Pain. *Pain Pract.* 2010;10:560-579.

4. Fujii K, Yamazaki M, Kang JD, et al. Discogenic Back Pain: Literature Review of Definition, Diagnosis, and Treatment. *JBMR Plus.* 2019;3:e10180.

5. Cheung KMC, Karppinen J, Chan D, et al. Prevalence and pattern of lumbar magnetic resonance imaging changes in a population study of one thousand forty-three individuals. *Spine (Phila Pa 1976).* 2009 Apr 20;34:934-940.

6. Schwarzer AC, Aprill CN, Derby R, Fortin J, Kine G. The relative contributions of the disc and zygapophyseal joint in chronic low back pain. *Spine (Phila Pa 1976).* 1994 Apr 1;19:801-806.

7. Verrills P, Nowesenitz G, Barnard A. Prevalence and Characteristics of Discogenic Pain in Tertiary Practice: 223 Consecutive Cases Utilizing Lumbar Discography. *Pain Med (United States).* 2015;16:1490-1499.

8. DePalma MJ, Ketchum JM, Saullo T. What Is the Source of Chronic Low Back Pain and Does Age Play a Role? *Pain Med.* 2011;12:224-233.

9. Manchikanti L. Evaluation of the Relative Contributions of Various Structures in ChronicLow Back Pain. *Pain Physician.* 2001;4:308-316.

10. Vo NV., Hartman RA, Patil PR, et al. Molecular mechanisms of biological aging in intervertebral discs. *J Orthop Res.* 2016;34:1289-306.

11. Singh K, Masuda K, Thonar EJMA, An HS, Cs-Szabo G. Age-related changes in the extracellular matrix of nucleus pulposus and anulus fibrosus of human intervertebral disc. *Spine (Phila Pa*

- 1976). 2009;34:10-16.
12. Patil P, Niedernhofer LJ, Robbins PD, Lee J, Sowa G, Vo N. Cellular Senescence in Intervertebral Disc Aging and Degeneration. *Curr Mol Biol Reports*. 2018;4:180-190.
13. Kos N, Gradisnik L, Velnar T. A Brief Review of the Degenerative Intervertebral Disc Disease. *Med Arch (Sarajevo, Bosnia Herzegovina)*. 2019;73:421-424.
14. Bibby SRS, Jones DA, Ripley RM, Urban JPG. Metabolism of the intervertebral disc: Effects of low levels of oxygen, glucose, and pH on rates of energy metabolism of bovine nucleus pulposus cells. *Spine (Phila Pa 1976)*. 2005;30:487-496.
15. Liang CZ, Li H, Tao YQ, et al. The relationship between low pH in intervertebral discs and low back pain: A systematic review. *Arch Med Sci*. 2012;8:952-956.
16. Keshari KR, Lotz JC, Link TM, Hu S, Majumdar S, Kurhanewicz J. Lactic acid and proteoglycans as metabolic markers for discogenic back pain. *Spine (Phila Pa 1976)*. 2008;33:312-317.
17. Schwarzer AC, Aprill CN, Derby R, Fortin J, Kine G, Bogduk N. The prevalence and clinical features of internal disc disruption in patients with chronic low back pain. *Spine (Phila Pa 1976)*. 1995;20:1878-1883.
18. Brinjikji W, Luetmer PH, Comstock B, et al. Systematic literature review of imaging features of spinal degeneration in asymptomatic populations. *Am J Neuroradiol*. 2015;36:811-816.
19. Manchikanti L, Benyamin RM, Singh V, et al. An update of the systematic appraisal of the accuracy and utility of lumbar discography in chronic low back pain. *Pain Physician*. 2013 ;16(2 Suppl):SE55-95
20. Laslett M, Oberg B, Aprill CN, et al. Centralization as a predictor of provocation discography results in chronic low back pain, and the influence of disability and distress on diagnostic power. *Spine J*. 2005;5:370-80.
21. Laslett M, Aprill CN, McDonald B, Oberg B. Clinical predictors of lumbar provocation discography: a study of clinical predictors of lumbar provocation discography. *Eur Spine J*. 2006;15(:1473-84

- 1  
2  
3 1 22. Manchikanti L, Benyamin RM, Singh V, et al. An update of the systematic appraisal of the  
4  
5 2 accuracy and utility of lumbar discography in chronic low back pain. *Pain Physician*. 2013 ;16(2  
6  
7 3 Suppl):SE55-95.  
8  
9  
10 4 23. Wolfer LR, Derby R, Lee JE, Lee SH. Systematic review of lumbar provocation discography in  
11  
12 5 asymptomatic subjects with a meta-analysis of false-positive rates. *Pain Physician*.  
13  
14 6 2008;11:513-538.  
15  
16 7 24. McCormick ZL, DeFrancesch F, Loomba V, et al. Diagnostic Value, Prognostic Value, and Safety  
17  
18 8 of Provocation Discography. *Pain Med*. 2018 Jan 1;19:3-8.  
19  
20 9 25. Carragee EJ, Don AS, Hurwitz EL, et al. 2009 ISSLS prize winner: Does discography cause  
21  
22 10 accelerated progression of degeneration changes in the lumbar disc: A ten-year matched  
23  
24 11 cohort study. *Spine (Phila Pa 1976)*. 2009;34:2338-2345.  
25  
26 12 26. Cuellar JM, Stauff MP, Herzog RJ, et al. Does provocative discography cause clinically important  
27  
28 13 injury to the lumbar intervertebral disc? A 10-year matched cohort study. *Spine J*.  
29  
30 14 2016;16:273-280  
31  
32 15 27. Koetsier E, Kallewaard JW, Maino P. Commentary on: Does provocative discography cause  
33  
34 16 clinically important injury to the lumbar intervertebral disc? A 10-year matched cohort study,  
35  
36 17 by Cueller et al., *Spine J* 2016. *Spine J*. 2017;17:610-611.  
37  
38 18 28. McCormick ZL, Lehman VT, Plastaras CT, et al. Low-Pressure Lumbar Provocation Discography  
39  
40 19 According to Spine Intervention Society/International Association for the Study of Pain  
41  
42 20 Standards Does Not Cause Acceleration of Disc Degeneration in Patients with Symptomatic  
43  
44 21 Low Back Pain: A 7-Year Matched Coh. *Spine (Phila Pa 1976)*. 2019;44:E1161-E1168.  
45  
46 22 29. Van Middelkoop M, Rubinstein SM, Kuijpers T, et al. A systematic review on the effectiveness  
47  
48 23 of physical and rehabilitation interventions for chronic non-specific low back pain. *Eur Spine J*.  
49  
50 24 Published online 2011. 2011;20:19-39.  
51  
52 25 30. Kallewaard JW, Wintraecken VM, Geurts JW, et al. A multicenter randomized controlled trial  
53  
54 26 on the efficacy of intradiscal methylene blue injection for chronic discogenic low back pain:  
55  
56  
57  
58  
59  
60

- 1 The IMBI study. *Pain*. 2019;160:945-953.
- 2
- 3 1
- 4
- 5 2 31. Helm S, Hayek SM, Benyamin R, Manchikanti L. Systematic review of the effectiveness of
- 6
- 7 3
- 8 thermal annular procedures in treating discogenic low back pain. *Pain Physician*. 2009;12:207-
- 9
- 10 4
- 11 232.
- 12 5 32. Freeman BJC. IDET: A critical appraisal of the evidence. In: *European Spine Journal*. 2006;15
- 13
- 14 6
- 15 Suppl 3(Suppl 3):S448-457.
- 16 7 32. Huygen F, Kallewaard JW, van Tulder M, et al. "Evidence-Based Interventional Pain Medicine
- 17
- 18 According to Clinical Diagnoses": Update 2018. *Pain Pract*. 2019 Jul;19:664-675.
- 19 8
- 20 9 34. Pan A, Hai Y, Yang J, et al. Adjacent segment degeneration after lumbar spinal fusion compared
- 21
- 22 10
- 23 with motion-preservation procedures: a meta-analysis. *Eur Spine J*. Published online 2016.
- 24
- 25 11
- 26 2016;25:1522-1532.
- 27 12 35. Bydon M, De La Garza-Ramos R, Macki M, et al. Lumbar fusion versus nonoperative
- 28
- 29 13
- 30 management for treatment of discogenic low back pain: A systematic review and meta-
- 31
- 32 14
- 33 analysis of randomized controlled trials. *J Spinal Disord Tech*. 2014;27:297-304.
- 34 15 36. Wang X, Wanyan P, Tian JH, et al. Meta-analysis of randomized trials comparing fusion surgery
- 35
- 36 16
- 37 to non-surgical treatment for discogenic chronic low back pain. *J Back Musculoskelet Rehabil*.
- 38
- 39 17
- 40 2015;28:621-627.
- 41 18 37. Vallejo R, Gupta A, Cedeno DL, et al. Clinical Effectiveness and Mechanism of Action of Spinal
- 42
- 43 19
- 44 Cord Stimulation for Treating Chronic Low Back and Lower Extremity Pain: a Systematic
- 45
- 46 20
- 47 Review. *Curr Pain Headache Rep*. 2020;24:70.
- 48 21 38. Kallewaard JW, Edelbroek C, Terheggen M, et al. A Prospective Study of Dorsal Root Ganglion
- 49
- 50 22
- 51 Stimulation for Non-Operated Discogenic Low Back Pain. *Neuromodulation*. 2020;23:196-202.
- 52 23 39. Desai MJ, Mansfield JT, Robinson DM, et al. Regenerative Medicine for Axial and Radicular
- 53
- 54 24
- 55 Spine-Related Pain: A Narrative Review. *Pain Pract*. 2020;20:437-453.
- 56 25 40. Gullbrand SE, Schaer TP, Agarwal P, et al. Translation of an injectable triple-interpenetrating-
- 57
- 58 network hydrogel for intervertebral disc regeneration in a goat model. *Acta Biomater*.
- 59
- 60

- 1
- 2
- 3 1 2017;60:201-209.
- 4
- 5 2 41. Gan Y, Li P, Wang L, et al. An interpenetrating network-strengthened and toughened hydrogel
- 6
- 7 3 that supports cell-based nucleus pulposus regeneration. *Biomaterials*. 2017;136:12-28.
- 8
- 9
- 10 4 42. Yue JJ, Morgenstern R, Morgenstern C, et al. Shape memory hydrogels - A novel material for
- 11
- 12 5 treating age-related degenerative conditions of the Spine. *Eur Musculoskelet Rev*. 2011;6:184-
- 13
- 14 6 188.
- 15
- 16 7 43. Sing SH. Lumbar intradiscal treatments: Early evaluation of patient satisfaction in disc repair by
- 17
- 18 8 nucleus augmentation (GelStix™). *J Tissue Eng Regen Med*. 2012;6 45.
- 19
- 20
- 21 9 44. Ceylan A, Aşık İ, Özgencil GE, Erken B. Clinical results of intradiscal hydrogel administration
- 22
- 23 10 (GelStix) in lumbar degenerative disc disease. *Turkish J Med Sci*. 2019;49:1634-1639.
- 24
- 25
- 26 11 45. Pfirrmann CW, Metzdorf A, Zanetti M, et al. Magnetic resonance classification of lumbar
- 27
- 28 12 intervertebral disc degeneration. *Spine (Phila Pa 1976)*. 2001;26:1873-1878.
- 29
- 30 13 46. Adams MA, Dolan P, Hutton WC. The stages of disc degeneration as revealed by discograms. *J*
- 31
- 32 14 *Bone Jt Surg - Ser B*. 1986;68:36-41.
- 33
- 34 15 47. Merskey H, Bogduk N. Classification of Chronic Pain. Descriptions of Chronic Pain Syndromes
- 35
- 36 16 and Definitions of Pain Terms. 2nd ed. Seattle, WA: IASP Press; 1994.
- 37
- 38
- 39 17 48. Fukui S, Nitta K, Iwashita N, et al. Percutaneous intradiscal high-pressure injection of saline and
- 40
- 41 18 lidocaine in patients with lumbar intervertebral disc extrusion. *J Anesth*. 2012;26:786-789.
- 42
- 43
- 44 19 49. Cao P, Jiang L, Zhuang C, et al. Intradiscal injection therapy for degenerative chronic discogenic
- 45
- 46 20 low back pain with end plate Modic changes. *Spine J*. 2011;11:100-106.
- 47
- 48 21 50. Khot A, Bowditch M, Powell J, et al. The Use of Intradiscal Steroid Therapy for Lumbar Spinal
- 49
- 50 22 Discogenic Pain: A Randomized Controlled Trial. *Spine (Phila Pa 1976)*. 2004;29:833-836.
- 51
- 52
- 53 23 51. Dworkin RH, Turk DC, Farrar JT, et al. Core outcome measures for chronic pain clinical trials:
- 54
- 55 24 IMMPACT recommendations. *Pain*. 2005;113:9-19.
- 56
- 57 25 52. Fairbank JCT, Pynsent PB. The Oswestry Disability Index. *Spine (Phila Pa 1976)*. 2000;25:2940-
- 58
- 59 26 2953.
- 60

- 1  
2  
3 1 53. EuroQol--a new facility for the measurement of health-related quality of life. *Health Policy*.  
4  
5 2 1990;16:199-208.  
6  
7 3 54. Cheung JPY, Luk KDK. The relevance of high-intensity zones in degenerative disc disease. *Int*  
8  
9 4 *Orthop*. 2019;43:861-867.  
10  
11 5 55. Dudli S, Fields AJ, Samartzis D, et al. Pathobiology of Modic changes. *Eur Spine J*. 2016;25:3723-  
12  
13 6 3734.  
14  
15 7 56. Kyere KA, Than KD, Wang AC, et al. Schmorl's nodes. *Eur Spine J*. 2012;21:2115-2121.  
16  
17 8 57. Peters ML, Sommer M, Van Kleef M, et al. Predictors of physical and emotional recovery 6 and  
18  
19 9 12 months after surgery. *Br J Surg*. 2010;97:1518-1527.  
20  
21 10 58. Sullivan M, Bishop S, Pivik J. The pain catastrophizing scale: development and validation.  
22  
23 11 *Psychol Assess*. 1995;7:524-532.  
24  
25 12 59. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand*.  
26  
27 13 Published online 1983. 1983;67:361-370.  
28  
29 14 60. Nicholas MK. The pain self-efficacy questionnaire: Taking pain into account. *Eur J Pain*.  
30  
31 15 2007;11:153-163.  
32  
33 16 61. Reid MC, Williams CS, Gill TM. The relationship between psychological factors and disabling  
34  
35 17 musculoskeletal pain in community-dwelling older persons. *J Am Geriatr Soc*. 2003;51:1092-  
36  
37 18 1098.  
38  
39 19 62. Denison E, Åsenlöf P, Lindberg P. Self-efficacy, fear avoidance, and pain intensity as predictors  
40  
41 20 of disability in subacute and chronic musculoskeletal pain patients in primary health care. *Pain*.  
42  
43 21 2004;111:245-252.  
44  
45 22 63. Kolahi J, Bang H, Park J. Towards a proposal for assessment of blinding success in clinical trials:  
46  
47 23 Up-to-date review. *Community Dent Oral Epidemiol*. 2009;37:477-484.  
48  
49 24 64. Hróbjartsson A, Emanuelsson F, Thomsen ASS, et al. Bias due to lack of patient blinding in  
50  
51 25 clinical trials. A systematic review of trials randomizing patients to blind and nonblind sub-  
52  
53 26 studies. *Int J Epidemiol*. 2014;43:1272-1283.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

65. Harris PA, Taylor R, Thielke R, et al. Research electronic data capture (REDCap)-A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform.* 2009;42:377-381

66. Kahan BC, Morris TP. Reporting and analysis of trials using stratified randomisation in leading medical journals: review and reanalysis. *BMJ.* 2012;345:e5840-e5840.

67. Manchikanti L, Knezevic NN, Sanapati J, et al. Is Epidural Injection of Sodium Chloride Solution a True Placebo or an Active Control Agent? A Systematic Review and Meta-Analysis. *Pain Physician.* 2021;24:41-59.

**Acknowledgements**

Prof. Roberto S.G.M. Perez had an important role in the initiation of this study and helped substantially with the conception and design of the study. To our greatest regret, he passed away on 07.09.2017.

**Author statement**

EK, PM, JWK, LS, AC and SK designed the study. EK, PM, JWK, JD, LS, AC, PS, DK will conduct the study including patient recruitment and data collection. SK will conduct the data analysis and will conduct the interpretation of the data. EK drafted the manuscript with important intellectual input from JWK, PM, SK, AC, JD, MH, LS, PS, and DK. All authors approved the final manuscript. EK, JWK, PM, and JD will have complete access to the study data.

1  
2  
3 1 **Figure legends**  
4

5 2 **Figure 1**  
6  
7

8 3 1835S GelStix™. From left to right: 18 Gauge Needle, GelStix: dry, after 15 minutes hydration, after  
9  
10 4 30 minutes hydration, after 45 minutes hydration  
11

12 5  
13  
14 6 **Figure 2**  
15

16 7 Study flow chart  
17  
18

19 8  
20

21 9 **Figure 3**  
22

23 10 3A) Using fluoroscopic guidance, the needle is introduced using a standard posterolateral discography  
24  
25 11 approach. 3B) The protective cap is removed from the preloaded implant holder. 3C) The implant  
26  
27 12 holder is threaded onto the proximal end of the introducer needle. 3D) The holder stylet is pushed so  
28  
29 13 that the implant is driven completely into the introducer needle. 3E) The implant holder is removed.  
30  
31 14 The needle stylet is driven through the needle and bottomed out to deliver the GelStix™ completely  
32  
33 15 into the nucleus, keeping the needle tip centered in the nucleus. 3F) The needle tip will keep  
34  
35 16 centered approximately in the nucleus and the procedure will be repeated to insert additional  
36  
37 17 GelStix™.  
38  
39  
40

41 18  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60







Figure 1  
1835S GelStix™. From left to right: 18 Gauge Needle, GelStix: dry, after 15 minutes hydration, after 30 minutes hydration, after 45 minutes hydration  
694x517mm (72 x 72 DPI)






Figure 3

3A) Using fluoroscopic guidance, the needle is introduced using a standard posterolateral discography approach. 3B) The protective cap is removed from the preloaded implant holder. 3C) The implant holder is threaded onto the proximal end of the introducer needle. 3D) The holder stylet is pushed so that the implant is driven completely into the introducer needle. 3E) The implant holder is removed. The needle stylet is driven through the needle and bottomed out to deliver the GelStix™ completely into the nucleus, keeping the needle tip centered in the nucleus. 3F) The needle tip will keep centered approximately in the nucleus and the procedure will be repeated to insert additional GelStix™.

338x190mm (96 x 96 DPI)

# Reporting checklist for protocol of a clinical trial.

Based on the SPIRIT guidelines.

## Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the SPIRIT reporting guidelines, and cite them as:

Chan A-W, Tetzlaff JM, Gøtzsche PC, Altman DG, Mann H, Berlin J, Dickersin K, Hróbjartsson A, Schulz KF, Parulekar WR, Krleža-Jerić K, Laupacis A, Moher D. SPIRIT 2013 Explanation and Elaboration: Guidance for protocols of clinical trials. BMJ. 2013;346:e7586

			Page
Reporting Item			Number
<b>Administrative information</b>			
Title	<a href="#">#1</a>	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	1

1	Trial registration	<a href="#">#2a</a>	Trial identifier and registry name. If not yet registered,	3
2			name of intended registry	
3				
4				
5				
6	Trial registration:	<a href="#">#2b</a>	All items from the World Health Organization Trial	NA
7			Registration Data Set	
8	data set			
9				
10				
11				
12	Protocol version	<a href="#">#3</a>	Date and version identifier	3
13				
14				
15	Funding	<a href="#">#4</a>	Sources and types of financial, material, and other	19
16			support	
17				
18				
19				
20	Roles and	<a href="#">#5a</a>	Names, affiliations, and roles of protocol contributors	1, 27
21				
22	responsibilities:			
23				
24	contributorship			
25				
26				
27				
28	Roles and	<a href="#">#5b</a>	Name and contact information for the trial sponsor	2
29				
30	responsibilities:			
31				
32	sponsor contact			
33				
34	information			
35				
36				
37				
38	Roles and	<a href="#">#5c</a>	Role of study sponsor and funders, if any, in study	2
39			design; collection, management, analysis, and	
40	responsibilities:		interpretation of data; writing of the report; and the	
41			decision to submit the report for publication, including	
42	sponsor and funder		whether they will have ultimate authority over any of	
43			these activities	
44				
45				
46				
47				
48				
49				
50				
51				
52	Roles and	<a href="#">#5d</a>	Composition, roles, and responsibilities of the	NA
53			coordinating centre, steering committee, endpoint	
54	responsibilities:		adjudication committee, data management team, and	
55				
56	committees			
57				
58				
59				
60				

other individuals or groups overseeing the trial, if  
applicable (see Item 21a for data monitoring committee)

## Introduction

Background and rationale	<a href="#">#6a</a>	Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention	5-7
Background and rationale: choice of comparators	<a href="#">#6b</a>	Explanation for choice of comparators	5-7
Objectives	<a href="#">#7</a>	Specific objectives or hypotheses	7
Trial design	<a href="#">#8</a>	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, non-inferiority, exploratory)	7
<b>Methods:</b>			
<b>Participants, interventions, and outcomes</b>			
Study setting	<a href="#">#9</a>	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will be collected. Reference to where list of study sites can be obtained	8

1	Eligibility criteria	<a href="#">#10</a>	Inclusion and exclusion criteria for participants. If	8-10
2				
3			applicable, eligibility criteria for study centres and	
4			individuals who will perform the interventions (eg,	
5			surgeons, psychotherapists)	
6				
7				
8				
9				
10				
11	Interventions:	<a href="#">#11a</a>	Interventions for each group with sufficient detail to allow	10-11
12				
13	description		replication, including how and when they will be	
14			administered	
15				
16				
17				
18				
19	Interventions:	<a href="#">#11b</a>	Criteria for discontinuing or modifying allocated	16
20				
21	modifications		interventions for a given trial participant (eg, drug dose	
22			change in response to harms, participant request, or	
23			improving / worsening disease)	
24				
25				
26				
27				
28				
29	Interventions:	<a href="#">#11c</a>	Strategies to improve adherence to intervention	NA
30				
31	adherence		protocols, and any procedures for monitoring adherence	
32			(eg, drug tablet return; laboratory tests)	
33				
34				
35				
36	Interventions:	<a href="#">#11d</a>	Relevant concomitant care and interventions that are	11
37				
38	concomitant care		permitted or prohibited during the trial	
39				
40				
41				
42	Outcomes	<a href="#">#12</a>	Primary, secondary, and other outcomes, including the	12-14
43				
44			specific measurement variable (eg, systolic blood	
45			pressure), analysis metric (eg, change from baseline,	
46			final value, time to event), method of aggregation (eg,	
47			median, proportion), and time point for each outcome.	
48				
49			Explanation of the clinical relevance of chosen efficacy	
50			and harm outcomes is strongly recommended	
51				
52				
53				
54				
55				
56				
57				
58				
59				
60				

Participant timeline	<a href="#">#13</a>	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure)	12, Fig. 2
Sample size	<a href="#">#14</a>	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	14
Recruitment	<a href="#">#15</a>	Strategies for achieving adequate participant enrolment to reach target sample size	8, 17
<b>Methods:</b>			
<b>Assignment of interventions (for controlled trials)</b>			
Allocation: sequence generation	<a href="#">#16a</a>	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	14
Allocation concealment mechanism	<a href="#">#16b</a>	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque,	14-15



1		sealed envelopes), describing any steps to conceal the	
2			
3		sequence until interventions are assigned	
4			
5			
6	Allocation:	<a href="#">#16c</a> Who will generate the allocation sequence, who will	14-15
7			
8	implementation	enrol participants, and who will assign participants to	
9			
10		interventions	
11			
12			
13	Blinding (masking)	<a href="#">#17a</a> Who will be blinded after assignment to interventions	14-15
14			
15		(eg, trial participants, care providers, outcome	
16			
17		assessors, data analysts), and how	
18			
19			
20			
21	Blinding (masking):	<a href="#">#17b</a> If blinded, circumstances under which unblinding is	15
22			
23	emergency	permissible, and procedure for revealing a participant's	
24			
25	unblinding	allocated intervention during the trial	
26			
27			
28			
29	<b>Methods: Data</b>		
30			
31	collection,		
32			
33	management, and		
34			
35	analysis		
36			
37			
38			
39	Data collection plan	<a href="#">#18a</a> Plans for assessment and collection of outcome,	15
40			
41		baseline, and other trial data, including any related	
42			
43		processes to promote data quality (eg, duplicate	
44			
45		measurements, training of assessors) and a description	
46			
47		of study instruments (eg, questionnaires, laboratory	
48			
49		tests) along with their reliability and validity, if known.	
50			
51			
52		Reference to where data collection forms can be found,	
53			
54		if not in the protocol	
55			
56			
57			
58			
59			
60			

1	Data collection plan:	<a href="#">#18b</a>	Plans to promote participant retention and complete	15
2				
3	retention		follow-up, including list of any outcome data to be	
4			collected for participants who discontinue or deviate from	
5			intervention protocols	
6				
7				
8				
9				
10				
11	Data management	<a href="#">#19</a>	Plans for data entry, coding, security, and storage,	15
12			including any related processes to promote data quality	
13			(eg, double data entry; range checks for data values).	
14			Reference to where details of data management	
15			procedures can be found, if not in the protocol	
16				
17				
18	Statistics: outcomes	<a href="#">#20a</a>	Statistical methods for analysing primary and secondary	16-17
19			outcomes. Reference to where other details of the	
20			statistical analysis plan can be found, if not in the	
21			protocol	
22				
23				
24	Statistics: additional	<a href="#">#20b</a>	Methods for any additional analyses (eg, subgroup and	16-17
25	analyses		adjusted analyses)	
26				
27				
28	Statistics: analysis	<a href="#">#20c</a>	Definition of analysis population relating to protocol non-	16
29	population and		adherence (eg, as randomised analysis), and any	
30	missing data		statistical methods to handle missing data (eg, multiple	
31			imputation)	
32				
33				
34	<b>Methods: Monitoring</b>			
35				
36				
37				
38				
39	Data monitoring:	<a href="#">#21a</a>	Composition of data monitoring committee (DMC);	17
40				
41	formal committee		summary of its role and reporting structure; statement of	
42			whether it is independent from the sponsor and	
43				
44				
45				
46				
47				
48				
49				
50				
51				
52				
53				
54				
55				
56				
57				
58				
59				
60				

1		competing interests; and reference to where further	
2		details about its charter can be found, if not in the	
3		protocol. Alternatively, an explanation of why a DMC is	
4		not needed	
5			
6			
7			
8			
9			
10	Data monitoring:	<a href="#">#21b</a> Description of any interim analyses and stopping	17
11	interim analysis	guidelines, including who will have access to these	
12		interim results and make the final decision to terminate	
13		the trial	
14			
15			
16			
17			
18			
19			
20	Harms	<a href="#">#22</a> Plans for collecting, assessing, reporting, and managing	14
21		solicited and spontaneously reported adverse events	
22		and other unintended effects of trial interventions or trial	
23		conduct	
24			
25			
26			
27			
28			
29			
30	Auditing	<a href="#">#23</a> Frequency and procedures for auditing trial conduct, if	NA
31		any, and whether the process will be independent from	
32		investigators and the sponsor	
33			
34			
35			
36			
37			
38	Ethics and		
39	dissemination		
40			
41			
42			
43	Research ethics	<a href="#">#24</a> Plans for seeking research ethics committee /	19
44	approval	institutional review board (REC / IRB) approval	
45			
46			
47			
48	Protocol	<a href="#">#25</a> Plans for communicating important protocol	19
49	amendments	modifications (eg, changes to eligibility criteria,	
50		outcomes, analyses) to relevant parties (eg,	
51		investigators, REC / IRBs, trial participants, trial	
52		registries, journals, regulators)	
53			
54			
55			
56			
57			
58			
59			
60			

Consent or assent	<a href="#">#26a</a>	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	19
Consent or assent: ancillary studies	<a href="#">#26b</a>	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	NA
Confidentiality	<a href="#">#27</a>	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	19
Declaration of interests	<a href="#">#28</a>	Financial and other competing interests for principal investigators for the overall trial and each study site	19
Data access	<a href="#">#29</a>	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	27
Ancillary and post trial care	<a href="#">#30</a>	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	NA
Dissemination policy: trial results	<a href="#">#31a</a>	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	15-16

1	Dissemination policy: <a href="#">#31b</a>	Authorship eligibility guidelines and any intended use of	NA
2			
3	authorship	professional writers	
4			
5			
6	Dissemination policy: <a href="#">#31c</a>	Plans, if any, for granting public access to the full	NA
7			
8	reproducible	protocol, participant-level dataset, and statistical code	
9			
10	research		
11			
12			
13			

14 **Appendices**

15			
16			
17	Informed consent	<a href="#">#32</a> Model consent form and other related documentation	appendix
18			
19	materials	given to participants and authorised surrogates	1
20			
21			
22	Biological specimens	<a href="#">#33</a> Plans for collection, laboratory evaluation, and storage of	NA
23			
24		biological specimens for genetic or molecular analysis in	
25			
26		the current trial and for future use in ancillary studies, if	
27			
28		applicable	
29			
30			
31			

32 The SPIRIT Explanation and Elaboration paper is distributed under the terms of the Creative  
33 Commons Attribution License CC-BY-NC. This checklist was completed on 20. May 2021 using  
34 <https://www.goodreports.org/>, a tool made by the [EQUATOR Network](#) in collaboration with  
35 [Penelope.ai](#)  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60